

# Inventory and world geographical distribution of the olive tree (*Olea europaea* L.) diseases caused by viruses, bacteria and phytoplasma

Mohamed Chliyeh, Amina Ouazzani Touhami, Karima Selmaoui, Rachid Benkirane,  
Allal Douira\*

Laboratoire de Botanique, Biotechnologie et Protection des Plantes, Département de Biologie, Université Ibn Tofail, Faculté des Sciences, BP. 133, Kénitra, Maroc (Morocco).

**Abstract**— The objective of this work is to prepare an Inventory about the viral and bacterial diseases of the olive tree (*Olea europaea* L.) and to study their Geographical distribution in the world. Fifteen viruses were considered as pathogens on the olive tree, these viruses were reported in 22 countries in five continents showing different symptoms on different olive tree varieties. All viruses were reported in Italy, some of these viruses were found only in the European continent, such as Olive vein yellows associated virus (OVYaV), Olive yellow mottling and decline associated virus (OYMDaV), Olive mild mosaic virus (OMMV) and Olive semilatent virus (OSLV). Two bacteria were reported as pathogen on the olive tree; *Pseudomonas savastanoi*. pv. *savastanoi* responsible bacteria of the olive knot; a serious disease of olive (*Olea europaea* L.) in the Mediterranean region. This bacteria was detected in 22 countries in five continents, the majority of these countries exist in the Mediterranean Basin. Another bacteria was *Xylella fastidiosa* responsible of the quick decline syndrome of olive (OQDS), was detected in Europe in four countries (Italy, Spain, Portugal and Slovenia) and in America in three countries (USA, Argentina and Brazil). Phtoplasmas was reported in 5 countries in three continents (Europe, Asia and Australia).

**Keywords**— viruses, bacteria, phytoplasma, olive tree.

## I. INTRODUCTION

Oleaceae family or the family of dicotyledons includes 30 genera (Grohmann, 1981; Cronquist, 1981) of deciduous trees and shrubs including olive tree and its relatives numbering about 600 species (Bianco, 1990). The family is divided into several tribes, that is, Fontanesieae, Forsythiaeae, Jasmineae, Myxopyreae, and Oleaeae (USDA, 2003; Bartolini and Petruccelli, 2002). These are mostly native of all continents except the Antarctic, including tropical, subtropical, and temperate regions of the world (Grohmann, 1981; Wallander and Albert, 2000). Oleaceae

is best grown in Asia and Malaysia especially tropical and temperate regions of Asia (Perez *et al.*, 2005). The genus *Olea* got its name from the Greek “elaia” and the Latin “oleum,” but it is known by nearly 80 different names (Medai *et al.*, 2001). The *Olea* genus comprises 30 species (Bracci *et al.*, 2011) but *Olea europaea* L. is the most popular member of the *Olea* genus (Kaniewski *et al.*, 2012). It is the only species of this genus which is used as food (Sarwar, 2013) and is found in the Mediterranean region (Zohary *et al.*, 2012).

The Olive is an important crop in terms of both its commercial value and the role it plays in the rural economy of the Mediterranean region, with its millions of producers (Gonzalez-Andujar, 2009). Olive tree and its fruit are also important in context of religion. Olives are narrated several times in the Bible, both in the New and Old Testaments (Ryan and Robard, 1998). Olive has also been praised as a blessed tree and fruit in the Holy Quran (Quran, Chapter 24 Al-Noor, Verse 35). Olives are not used as a natural fruit because of their extremely bitter taste but are rather consumed either as olive oil or table olives (Kanakis *et al.*, 2013). Olive oil market is very significant in the olive industry as approximately 90% of annually produced olives go for oil processing (Sibbett *et al.*, 2005).

The olive tree is one of the oldest cultivated trees on the planet earth (Liphshitz *et al.*, 1991). The cultivation of olive started in ancient times and it dates back more than 7000 years. According to archaeological reports olives were cultivated for commercial purposes in Crete in 3000 BC by the Minoan civilization. The use of olive oil for body health can be found in ancient Greek literature (Gooch, 2005).

The therapeutic utilities of *O. europaea* have been indicated in traditional medicine. It has been known to reduce blood sugar, cholesterol, and uric acid. It has also been used to treat diabetes, hypertension, inflammation, diarrhea, respiratory and urinary tract infections, stomach

and intestinal diseases, asthma, hemorrhoids, rheumatism, laxative, mouth cleanser, and as a vasodilator (Hashmi *et al.*, 2015).

Unfortunately, olive is subjected to be attacked with a variety of pathogens, which affect its health, yield and its oil quality (Sergeeva *et al.*, 2005; Sanei *et al.*, 2004). Several biotic and abiotic factors affect olive tree consequently reducing the tree health and vigour, crop yield and quality of the extracted oil (Sanei *et al.*, 2012; Rhouma *et al.*, 2010; Palumbo *et al.*, 2006; Gutierrez *et al.*, 1999; Kiritsakis, 1998). Olive cultivar is one of the most important factors that affect the quality of olive oil<sup>34</sup>. In addition to the effects of cultivar, the degree of fruit ripeness, and the industrial processes used for oil extraction, as well as the quality of the fruit from which oil is extracted has a great effect on olive oil quality (Garcia and Yousfi, 2006). The establishment of new olive plantations, the incidence of wilting, dieback and death of young trees has also increased, alarming olive farmers (Chliyeh *et al.*, 2014).

Phytoplasmas constitute a monophyletic clade within the class Mollicutes. Their classification has been possible by the use of restriction fragment length polymorphism analysis (RFLP) and sequencing of the conserved 16S rRNA gene; the majority of phytoplasmas were classified into 15 (Lee *et al.*, 1998; Montano *et al.*, 2001) or 20 16S rDNA groups, encompassing a large number of subgroups (Caglayan *et al.*, 2011) On this basis, the phytoplasmas identified in olive trees have been classified as 16S-IB (Aster yellows group), 16S-VA (Elm yellows group) and 16S-XIIA (Stolbur group) (Pasquini *et al.*, 2000).

Some viruses are well characterized and infect many crops whereas others are apparently restricted to olive trees. Some of them are rare and have been detected sporadically while others are present in different countries and have a high incidence of infection in olive plants (Çağlayan *et al.*, 2011). For the sake of presentation, these viruses can be divided in two main groups: viruses identified for the first time in olive trees (viruses named after olives) and viruses already known to infect different crops and subsequently identified in olive trees (other viruses) (Çağlayan *et al.*, 2011). As the same, the olive tree is susceptible to different bacteria which can provoke damage and yield loss.

The aim of this work is to prepare an Inventory about the viral and bacterial diseases of the olive tree (*Olea europaea* L.) and to study their Geographical distribution in the world.

## II. CATALOG

### II-1- viruses:

**• Olive leaf yellowing associated virus (OLYaV): Croitia** (Godena *et al.*, 2016; Luigi *et al.*, 2011; Bjelis *et al.*,

*al.*, 2007); **Italy** (Loconsole *et al.*, 2013; Cutuli *et al.*, 2011; Rwahnih *et al.*, 2011; Savino *et al.*, 2011; Al-abdallah *et al.*, 2010; Montemurro *et al.*, 2008; El Beaino *et al.*, 2007, 2005; Fagioli *et al.*, 2005; Savino *et al.*, 2004; Saponari *et al.*, 2004, 2002a, 2001a, 2001b; Albanese *et al.*, 2003; Martelli, 2002; Bellini *et al.*, 2002; Bartolini *et al.*, 2002; Bottalico *et al.*, 2002; Abou Ghanem *et al.*, 2001; Savino *et al.*, 1996; Grieco *et al.*, 2000; Sabanadzovic *et al.*, 1999; Maretti and Gallitelli, 1985); **California** (Essakhi *et al.*, 2006; Al Rwahnih *et al.*, 2011); **Albania** (Luigi *et al.*, 2009); **Turkey** (Erglmez and Erkan, 2016, 2014; Çgfcg and Serce, 2014); **Tunisia** (Boulila, 2016; Rwahnih *et al.*, 2011; El Air *et al.*, 2011); **Lebanon** (Fadel *et al.*, 2005); **Egypt** (Rwahnih *et al.*, 2011; Youssef *et al.*, 2010); **Spain** (Rwahnih *et al.*, 2011; Bertolini *et al.*, 1998); **Israel** (Al Rwahnih *et al.*, 2011); **Greece** (Rwahnih *et al.*, 2011); **Australia** (Rwahnih *et al.*, 2011); **Cyprus** (Rwahnih *et al.*, 2011); **Chile** (Rwahnih *et al.*, 2011); **France** (Rwahnih *et al.*, 2011); **Syria** (Rwahnih *et al.*, 2011; Abdine *et al.*, 2007; Al Abdullah, 2004); **Palestine** (Rwahnih *et al.*, 2011); **Morocco** (Rwahnih *et al.*, 2011); **Algeria** (Rwahnih *et al.*, 2011).

**• Cherry leaf roll virus (CLRV): Croitia** (Godena *et al.*, 2016, 2012; Luigi *et al.*, 2011; Bjelis *et al.*, 2007); **Italy** (Ferretti *et al.*, 2017; Martelli, 2013a, 2013b; Loconsole *et al.*, 2013; Savino *et al.*, 2011; Al-abdallah *et al.*, 2010; Fagioli *et al.*, 2005; Savino *et al.*, 2004; Saponari *et al.*, 2004, 2002a, 2002b, 2002c, 2001a, 2001b; Martelli, 2002; Bartolini *et al.*, 2002; Bellini *et al.*, 2002; Bottalico *et al.*, 2002; Pantaleo *et al.*, 2001; Grieco *et al.*, 2000; Leitao *et al.*, 1999; Martelli *et al.*, 1995; Maretti and Gallitelli, 1985; Savino and Gallitelli, 1981); **Spain** (Bertolini *et al.*, 2001a, 2001b, 1998; Grieco *et al.*, 2000); **Portugal** (Felix *et al.*, 2002; Grieco *et al.*, 2000); da Clara Henriques, 1994; **Lebanon** (Fadel *et al.*, 2005); **Tunisia** (Boulila, 2016; El Air *et al.*, 2011); **Egypt** (Youssef *et al.*, 2010); **Turkey** (Erglmez and Erkan, 2016, 2014; Çgfcg and Serce, 2014; Yalçın *et al.*, 2008; Beler and Açıkgöz, 2005; Caglayan *et al.*, 2004); **Albania** (Luigi *et al.*, 2009); **Syria** (Abdine *et al.*, 2007; Al Abdullah *et al.*, 2005).

**• Strawberry latent ring spot virus (SLRSV): Croitia** (Godena *et al.*, 2016; Luigi *et al.*, 2011; Bjelis *et al.*, 2007); **Italy** (Loconsole *et al.*, 2013; Savino *et al.*, 2011; Fagioli *et al.*, 2005, 2002; Savino *et al.*, 2004; Pantaleo *et al.*, 2001; Ferretti *et al.*, 2002; Bellini *et al.*, 2002; Martelli, 2002; Saponari *et al.*, 2002a, 2002b, 2001a, 2001b; Bartolini *et al.*, 2002; Grieco *et al.*, 2000; Martelli *et al.*, 1995; Barba, 1993; Marte *et al.*, 1986; Maretti and Gallitelli, 1985; Savino *et al.*, 1979; Pacini and Cresti, 1977); **Spain** (Bertolini *et al.*, 2001a; 2001b, 1998; da Clara Henriques, 1994; Grieco *et al.*, 2000); **Portugal**

(Félix *et al.*, 2002; Grieco *et al.*, 2000; Leitao *et al.*, 1999; Henriques *et al.*, 1992, 1990); **Lebanon** (Fadel *et al.*, 2005); **Tunisia** (Boulila, 2016; El Air *et al.*, 2011); **Egypt** (EL-MORSY *et al.*, 2017; Youssef *et al.*, 2010), **California** (Saponari and Savino, 2003); **Turkey** (Çğfçg and Serçe, 2016, 2014; Yalçın *et al.*, 2008; Beler and Açıkgöz, 2005; Caglayan *et al.*, 2004), **Syria** (Abdine *et al.*, 2007; Al Abdullah *et al.*, 2005, 2004); **Albania** (Luigi *et al.*, 2009).

• **Arabis mosaic virus (ArMV): Croitia** (Godena *et al.*, 2016; Luigi *et al.*, 2011); **Italy** (Loconsole *et al.*, 2013; Savino *et al.*, 2011; Al-abdallah *et al.*, 2010; Faggioli *et al.*, 2005; Savino *et al.*, 2004; Martelli, 2002; Bellini *et al.*, 2002; Bartolini *et al.*, 2002; Saponari *et al.*, 2002a, 2002b, 2001a, 2001b; Pantaleo *et al.*, 2001; Grieco *et al.*, 2000; Martelli *et al.*, 1995; Marette and Gallitelli, 1985; Savino *et al.*, 1979); **Spain** (Bertolini *et al.*, 2001a, 1998); **Lebanon** (Fadel *et al.*, 2005); **Tunisia** (Boulila, 2016; El Air *et al.*, 2011); **Portugal** (Félix *et al.*, 2002; da Clara Henriques, 1994; **Egypt** (Youssef *et al.*, 2010); **Turkey** (Erğlmez and Erkan, 2016; Çğfçg and Serçe, 2014; Yalçın *et al.*, 2008; Beler and Açıkgöz, 2005; Caglayan *et al.*, 2004); **Albania** (Spahiu and Veizi, 2012; Luigi *et al.*, 2009); **Croatia** (Bjelis *et al.*, 2007); **California** (Rwahnih *et al.*, 2011); **Syria** (Abdine *et al.*, 2007; Al Abdullah *et al.*, 2005; Al Abdullah, 2004).

• **Olive latent virus-1 (OLV-1): Italy** (Martelli, 2013a, 2013b, 2002; Loconsole *et al.*, 2013; Al-abdallah *et al.*, 2010; Pantaleo *et al.*, 2006, 2001, 1999; Castellano *et al.*, 2005; Savino *et al.*, 2004; Bellini *et al.*, 2002; Bartolini *et al.*, 2002; Bellini *et al.*, 2002; Saponari *et al.*, 2002a, 2002b, 2002c, 2001a, 2001b; Grieco *et al.*, 1996a, 1995; Merciega *et al.*, 1996; Savino *et al.*, 1996; Martelli *et al.*, 1995; Vanessa, 1995; Castellano *et al.*, 1987; Marette and Gallitelli, 1985; Gallitelli and Savino, 1985), **Jordan** (Grieco *et al.*, 2000; Merciega *et al.*, 1996; Martelli *et al.*, 1996, 1995); **Portugal** (Varanda *et al.*, 2015; 2014a, 2014c; 2010, 2009, 2008; dos Santos, 2015; Gaspar, 2014; Felix *et al.*, 2011, 2007, 2001; Varanda, 2011; Félix, 2007, 2006a, 2006b; 2005a, 2005b, 2004; Felix, 2007; Felix and Clara, 2006, 2000, 1998; Lobão *et al.*, 2002; Cardoso *et al.*, 2005; Grieco *et al.*, 2000; da Clara Henriques, 1994); **Lebanon** (Fadel *et al.*, 2005), **Syria** (Abdine *et al.*, 2007; Al Abdullah *et al.*, 2005, 2004); **Turkey** (Erğlmez and Erkan, 2014; Çğfçg and Serçe, 2014; Yalçın *et al.*, 2008; Serce *et al.*, 2007; Grieco *et al.*, 2000; Merciega *et al.*, 1996); **Japan** (Kanematsu *et al.*, 2001); **Spain** (Bertolini *et al.*, 2001a, 1998; Grieco *et al.*, 2000); **Tunisia** (Boulila, 2016; El Air *et al.*, 2011); **Egypt** (Youssef *et al.*, 2010); **Albania** (Luigi *et al.*, 2009); **Croatia** (Godena *et al.*, 2016; Luigi *et al.*, 2011; Bjelis *et al.*, 2007); **California** (Al Rwahnih *et al.*, 2011).

• **Cucumber mosaic virus (CMV): Croitia** (Godena *et al.*, 2016; Luigi *et al.*, 2011; Bjelis *et al.*, 2007); **Italy** (Loconsole *et al.*, 2013; Savino *et al.*, 2011; Al-abdallah *et al.*, 2010; Faggioli *et al.*, 2005; Martelli, 2002; Saponari *et al.*, 2002a, 2001a; Bellini *et al.*, 2002; Bartolini *et al.*, 2002; Grieco *et al.*, 2000; Martelli *et al.*, 1995; Marette and Gallitelli, 1985; Savino and Gallitelli, 1983a); **Spain** (Rwahnih *et al.*, 2011; Bertolini *et al.*, 2001a, 2001b, 1998; Grieco *et al.*, 2000); **Portugal** (Gaspar, 2014; Félix and Clara, 2006; Félix *et al.*, 2002; Grieco *et al.*, 2000; Leitao *et al.*, 1999; da Clara Henriques, 1994; Rei *et al.*, 1993); **Tunisia** (Boulila, 2016; Rwahnih *et al.*, 2011; El Air *et al.*, 2011); **Egypt** (Rwahnih *et al.*, 2011; Youssef *et al.*, 2010); **California** (Al Rwahnih *et al.*, 2011); **Albania** (Luigi *et al.*, 2009); **Turkey** (Erğlmez and Erkan, 2016, 2014; Çğfçg and Serçe, 2014; Beler and Açıkgöz, 2005; Caglayan *et al.*, 2004); **Syria** (Rwahnih *et al.*, 2011; Abdine *et al.*, 2007; Al Abdullah, 2004); **Algeria** (Rwahnih *et al.*, 2011); **Australia** (Rwahnih *et al.*, 2011); **France** (Rwahnih *et al.*, 2011); **Cyprus** (Rwahnih *et al.*, 2011); **Israel** (Rwahnih *et al.*, 2011); **Morocco** (Rwahnih *et al.*, 2011); **Palestine** (Rwahnih *et al.*, 2011); **Greece** (Rwahnih *et al.*, 2011); **Lebanon** (Choueiri *et al.*, 2015).

• **Olive latent virus-2 (OLV-2): Croitia** (Godena *et al.*, 2016; Luigi *et al.*, 2011; Bjelis *et al.*, 2007); **Italy** (Loconsole *et al.*, 2013; Savino *et al.*, 2011; Al-abdallah *et al.*, 2010; Faggioli *et al.*, 2005; Martelli, 2002; Bellini *et al.*, 2002; Bartolini *et al.*, 2002; Saponari *et al.*, 2002a, 2001a, 2001b; Pantaleo *et al.*, 2001; Grieco *et al.*, 1996b, 2002, 2000; Martelli *et al.*, 1995; Castellano *et al.*, 1987; Marette and Gallitelli, 1985; Savino *et al.*, 1984; Corte *et al.*, 1961); **Spain** (Bertolini *et al.*, 2001a, 1998; Grieco *et al.*, 2000); **Syria** (Abdine *et al.*, 2007; Al Abdullah *et al.*, 2005, 2004); **Lebanon** (Grieco *et al.*, 2000); **Tunisia** (Boulila, 2016; El Air *et al.*, 2011); **Egypt** (Youssef *et al.*, 2010); **Greece** (Vovlas *et al.*, 2002); **Turkey** (Erğlmez and Erkan, 2016, 2014; Çğfçg and Serçe, 2014; Yalçın *et al.*, 2008); **Albania** (Spahiu and Veizi, 2012; Luigi *et al.*, 2009); **Portugal** (Gaspar, 2014; Henriques *et al.*, 1992, 1990).

• **Olive latent virus-3: Syria** (Rwahnih *et al.*, 2011; Al Abdullah *et al.*, 2010); **Turkey** (Çğfçg and Serçe, 2016, 2014; Erğlmez and Erkan, 2014; Rwahnih *et al.*, 2011) Al Abdullah *et al.*, 2010); **Italy** (Rwahnih *et al.*, 2011; Elbeaino *et al.*, 2011; Al Abdullah *et al.*, 2010); **Lebanon** (Rwahnih *et al.*, 2011; Al Abdullah *et al.*, 2010); **Greece** (Rwahnih *et al.*, 2011; Al Abdullah *et al.*, 2010); **Malta** (Rwahnih *et al.*, 2011; Al Abdullah *et al.*, 2010); **Portugal** (Gaspar, 2014; Rwahnih *et al.*, 2011; Al Abdullah *et al.*, 2010); **Tunisia** (Rwahnih *et al.*, 2011; Al Abdullah *et al.*, 2010).

• **Olive semilatent virus (OSLV): Italy** (Loconsole *et al.*, 2013; 2010; Martelli *et al.*, 2013, 2002; Fagioli *et al.*, 2005; Bellini *et al.*, 2002; Bartolini *et al.*, 2002; Bertolini *et al.*, 2001b; Saponari *et al.*, 2001; Grieco *et al.*, 2000; Maretti and Gallitelli, 1985; Materazzi *et al.*, 1996); **Turkey** (Çğfçğ and Serçe, 2016, 2014); **Portugal** (Gaspar, 2014).

• **Olive mild mosaic virus (OMMV): Portugal** (Varanda *et al.*, 2015, 2014b, 2014c; 2011a, 2011b, 2010, 2009, 2008; dos Santos, 2015; Gaspar, 2014; Varanda *et al.*, 2014; Félix *et al.*, 2011, 2007; Cardoso *et al.*, 2012, 2009, 2005; Felix, 2007; Felix and Clara, 2006; Félix and Clara, 2002); **Italy** (Martelli, 2013a, 2013b; Loconsole *et al.*, 2013; Loconsole *et al.*, 2010; Bartolini *et al.*, 2002; Bellini *et al.*, 2002); **Greece** (Gratsia *et al.*, 2012); **Turkey** (Çğfçğ and Serçe, 2014); **Spain** (Bertolini *et al.*, 2001a).

• **Olive yellow mottling and decline associated virus (OYMDaV): Italy** (Martelli, 2002, 1999; Bartolini *et al.*, 2002; Saponari *et al.*, 2001; Grieco *et al.*, 2000; Savino *et al.*, 1996; Fagioli and Barba, 1995; Maretti and Gallitelli, 1985); **Turkey** (Çğfçğ and Serçe, 2014); **Portugal** (dos Santos, 2015; Gaspar, 2014); **Spain** (Bertolini *et al.*, 2001a).

• **Olive latent ringspot virus (OLRSV): Croitia** (Luigi *et al.*, 2011; Bjelis *et al.*, 2007); **Italy** (Loconsole *et al.*, 2013; Martelli *et al.*, 2013b; Al-abdallah *et al.*, 2010; Montemurro *et al.*, 2008; Fagioli *et al.*, 2005, 2002; Saponari *et al.*, 2002a; Martelli, 2002; Bellini *et al.*, 2002; Bartolini *et al.*, 2002; Pantaleo *et al.*, 2001; Saponari *et al.*, 2001; Alkowni *et al.*, 2000a, 2000b; Grieco *et al.*, 2000; Martelli *et al.*, 1995; Savino *et al.*, 1983; Savino and Gallitelli, 1983b); **Tunisia** (Boulila, 2016; El Air *et al.*, 2011); **Portugal** (Gaspar, 2014; Grieco *et al.*, 2000); **Egypt** (Youssef *et al.*, 2010); **Turkey** (Erglmez and Erkan, 2016, 2014; Çğfçğ and Serçe, 2014; Yalçın *et al.*, 2008; Caglayan *et al.*, 2004); **Syria** (Abdine *et al.*, 2007; Al Abdullah *et al.*, 2005, 2004); **Lebanon** (Choueiri *et al.*, 2015).

• **Olive vein yellows associated virus (OVYaV): Croitia** (Luigi *et al.*, 2011); **Italy** (Martelli, 2013; Bellini *et al.*, 2002; Bartolini *et al.*, 2002; Martelli *et al.*, 2002; Saponari *et al.*, 2001; Grieco *et al.*, 2000; Martelli, 1999; Martelli *et al.*, 1995; Fagioli and Barba, 1995; Maretti and Gallitelli, 1985); **Turkey** (Erglmez and Erkan, 2016, 2014; Çğfçğ and Serçe, 2014); **Portugal** (Gaspar, 2014, Henriques *et al.*, 1992, 1990).

• **Tobacco necrosis virus D (TNV): Croitia** (Godena *et al.*, 2016; Luigi *et al.*, 2011), **Portugal** (Varanda *et al.*, 2015a, 2015b, 2014a, 2014b, 2014c; dos Santos, 2015; Gaspar, 2014; Al-abdallah *et al.*, 2012; Varanda *et al.*, 2011; Félix *et al.*, 2011, 2007, 2005; Cardoso *et al.*, 2012; 2009, 2005, 2004; Varanda, 2011; Varanda *et al.*,

2010; Felix, 2007; Pantaleo *et al.*, 2006, 1999; Felix and Clara, 2006; 2002, 2000); **Italy** (Martelli, 2013a, 2013b, 2002; Loconsole *et al.*, 2013, 2010; Savino *et al.*, 2011; Bartolini *et al.*, 2002; Saponari *et al.*, 2001; Martelli *et al.*, 2002, 1996; Grieco *et al.*, 2000, 1992; Triolo *et al.*, 1996; Vanessa, 1995; Maretti and Gallitelli, 1985); **Albania** (Spahiu and Veizi, 2012; Luigi *et al.*, 2009); **California** (Rwahnih *et al.*, 2011); **Turkey** (Erglmez S., Erkan S., 2014; Çğfçğ and Serçe, 2014); **Spain** (Bertolini *et al.*, 2001a).

• **Tobacco mosaic virus (TMV): Italy** (Martelli, 2013a, 2013b, 2002; Loconsole *et al.*, 2013; Savino *et al.*, 2011; Loconsole *et al.*, 2010; Fagioli *et al.*, 2005; Bartolini *et al.*, 2002; Saponari *et al.*, 2001; Grieco *et al.*, 2000, 1992; Triolo *et al.*, 1996; Martelli *et al.*, 2002, 1996; Vanessa, 1995; Maretti and Gallitelli, 1985, Savino *et al.*, 1983); **California** (Rwahnih *et al.*, 2011); **Jordan** (Martelli *et al.*, 1995).

## II- 2- Bacteria:

• **Pseudomonas savastanoi. pv. Savastanoi :** **Spain** (Gómez-Lama *et al.*, 2017; Caballo-Ponce *et al.*, 2017; Pérez-Mendoza *et al.*, 2014; Passos da Silva *et al.*, 2014; Aragon *et al.*, 2015, 2014a; 2014b; Matas *et al.*, 2014, 2012, 2010, 2009; Maldonado-González *et al.*, 2013; Ramos *et al.*, 2012; Rodríguez-Moreno *et al.*, 2010, 2008; Pérez-Martínez *et al.*, 2010, 2008, 2007; Queseda *et al.*, 2012, 2010a, 2010b, 2007; Trapero *et al.*, 2009; Rodríguez-Moreno *et al.*, 2009, 2008; Cayuela *et al.*, 2006; Martos-Moreno *et al.*, 2006; Penyalver *et al.*, 2006, 2005, 2003, 2000, 1998; Rojas *et al.*, 2004; Bertolini, 2003; Bertolini *et al.*, 2003a, 2003b, 2002; García *et al.*, 2001; Lopez-Villalta, 1999; Rojas, 1999; Alvarez *et al.*, 1998; García de los Ríos, 1989; Jiménez Diaz, 1985; Beltra, 1961, 1958); **Italy** (Moretti *et al.*, 2017; 2016; Buonauro *et al.*, 2015; Moretti *et al.*, 2014a, 2014b, 2014c, 2014d, 2011; Passos da Silva *et al.*, 2014, 2013; Lamichhane and Varvaro, 2013; Marchi *et al.*, 2009, 2006, 2005; Hosni *et al.*, 2011, Tous *et al.*, 2010; Hosni, 2010; Tegli *et al.*, 2010; Sisto *et al.*, 2010, 2007, 2004, 2002, 2001; Cimmino *et al.*, 2008; Moretti *et al.*, 2008; Cimmino *et al.*, 2006; Catara *et al.*, 2005; Capuzzo *et al.*, 2005; Scorticini *et al.*, 2004; Cortese *et al.*, 2004; Cinelli *et al.*, 2014; 2003; Surico and Marchi, 2003; Martelli *et al.*, 2002; Morea *et al.*, 1999; Balestra and Varvaro, 1997; Surico, 1993, 1977; Lavermicocca *et al.*, 1999, 1998; Mugnai *et al.*, 2004; Varvaro and Martella, 1993; Tommasini, 1992; Surico and Iacobellis, 1991; Surico and Lavermicocca, 1989; Evidente and Surico, 1986; Casano, 1985; Surico *et al.*, 1985a, 1985b, 1984; Hassani *et al.*, 2003; Lavermicocca *et al.*, 2003, 2002; Godini eg al., 2002; Iacobellis, 2001; Sisto and Iacobellis, 1999, 1992; Iacobellis *et al.*, 1998 ; 1995, 1994, 1993; Capasso *et al.*, 1997, 1995; Proetta, 1995; Ercolani, 1999, 1985, 1983,

1978, 1970; Caponero *et al.*, 1995; Wells *et al.*, 1991; Graniti, 1993; Evidente *et al.*, 1986; Varvaro and Ferrulli, 1983; Varvaro and Surico, 1987; 1984; Baratta and Marco, 1981; Varvaro and Surico, 1978a, 1978b; Surico *et al.*, 1975; Botalico and Ercolani, 1971;; Sands *et al.*, 1970; Domenico, 1969; Ciccarone, 1950; Petri, 1915; Petri, 1909; Savastano, 1908, 1889, 1887, 1886, 1878); **France** (Rodríguez-Palenzuela *et al.*, 2010; Bardaji *et al.*, 2011; Gaudin *et al.*, 1994; Gardan *et al.*, 1992a; 1992b; Abu-Ghorrah, 1988); **Jordan** (Khlaif, 2006; Tehabsim *et al.*, 1991); **Tunisia** (Ghanney *et al.*, 2016; Trigui *et al.*, 2013; Krid *et al.*, 2012; 2011a, 2011b, 2010, 2009; Mehri *et al.*, 2009; Ouzari *et al.*, 2008; Jardak *et al.*, 2004; Boulila and Mahjoub, 1994; Boulila *et al.*, 1993); **California** (Nguyen *et al.*, 2017; Úrbez-Torres *et al.*, 2013; Fichtner *et al.*, 2012; Adaskaveg, *et.al.*, 2012; Teviotdale, 2005; Feil *et al.*, 2005; Teviotdale and Krueger, 2004, 1998; Rice *et al.*, 2003; Penalver *et al.*, 2000; Krueger *et al.*, 1999; Scroth, 1995; Teviotdale, 1994; Soby *et al.*, 1993; Wilson and Lindow, 1992; Da Costa E Silva and Kosuge, 1991; Janse, 1991, 1982; Gaffney *et al.*, 1990; Roberto *et al.*, 1990; Palm *et al.*, 1989; Glass and Kosuge, 1988, 1986; Roberto and Kosuge, 1987; Lang, 1987; Macdonald *et al.*, 1986; Martin, 1986; Yamada *et al.*, 1986; Smidt and Vidaver, 1986; Hutcheson and Kosuge, 1985; Comai and Kosuge, 1983; Comai *et al.*, 1982; Comai and Kosuge, 1982, 1980; Shoemaker *et al.*, 1979; Wilson and Ogawa, 1979; Smidt and Kosuge, 1978; Hartmann, 1973; Marlow and Kosuge, 1972; Wilson *et al.*, 1972; Kuo and Kosuge, 1969; Hutzinger and Kosuge. 1968; Kosuge *et al.*, 1966; Wilson, 1965, 1935; Wilson and Maggie, 1964, 1963, 1962; Hagen, 1963; Sutic and Dowson, 1963; Ark and Thompson, 1960; Hartmann, 1949; Hewitt, 1938; Brown, 1932; Smith, 1922, 1908; Horne *et al.*, 1912; Smith and Rorer, 1904); **Algeria** (Kacem *et al.*, 2009; Kacem and Kazouz, 2009); **Morocco** (Bouaichi *et al.*, 2015; Mahhou *et al.*, 2012; Benjama, 2003, 2002, 1994, 1990, 1989, 1988, 1987; Benjama *et al.*, 2002, 1992, 1993, 1987; Benjama and Gardan, 1995; Moukhli, 1984; Rieuf and Teasca, 1970); **Portugal** (Campos *et al.*, 2009; Gemas *et al.*, 2004; Oliveira and Luz 2003; Fernandes and Marcelo, 2002; Marcelo *et al.*, 1999); **New Zealand** (Young, 2014a, 2004b, 1987; Wilson *et al.*, 2004; Young *et al.*, 1996; Young and Triggs, 1994; Braithwaite *et al.*, 1999; Dye, 1956); **Lebanon** (Temsah *et al.*, 2008, 2007a, 2007b, 2007c; Saad and Hanna, 2002; Saad and Melkonian, 1992; Saad and Nienhaus, 1969; Weltzien, 1963); **Greece** (Hatzopoulos *et al.*, 2002; Roussos *et al.*, 2002; Roussos and Pontikis, 2002; Krueger *et al.*, 1999; Schroth, 1995; Pyrowolakis and Weltzin, 1974; Panagopoulos, 1993; Tjamos *et al.*, 1993; Schroth and Hildebrand, 1968; Schroth *et al.*, 1973, 1968); **Australia**

(Spooner-Hart *et al.*, 2009; Hall *et al.*, 2004, 2003; Kailis and Sweeney, 2002; Moffett, 1983; Sampson and Walker, 1982; Warcup and Talbot, 1981; Noble *et al.*, 1935; Adam and Pugsley, 1934 ); **Iraq** (Osman *et al.*, 1980a, 1980b); **Turkey** (Bozkurt *et al.*, 2014; Servi and Baştas, 2012; Mirik and Aysan, 2011; Kavak and Üstün, 2009; Mirik *et al.*, 2006; Basim *et al.*, 2006, 2000; Tatlı and Benlioğlu, 2004 ; Basim, 2002; Basim and Ersoy, 2001; Azeri, 1993); **Nepal** (Balestraa *et al.*, 2009); **Iran** (Rokni-Zadeh *et al.*, 2008); **Syria** (Alabdalla *et al.*, 2009); **China** (Haigen *et al.*, 2006); **Croatia** (Godena *et al.*, 2009; Škarica *et al.*, 1996; Žužić and Ciglar, 1987); **Egypt** (Ahmad *et al.*, 2009); **Japan** (Tsuji *et al.*, 2017; 2015).

• **Xylella fastidiosa** : **Italy** (Bleve *et al.*, 2016; Pashaei *et al.*, 2016; Mang *et al.*, 2016; Moussa *et al.*, 2016; Cornara *et al.*, 2016; Martelli, 2016, 2013; Loconsole *et al.*, 2016; 2014a; 2014b; Cornara *et al.*, 2016; 2014; Bosso *et al.*, 2016; Almeida, 2016; Saponari *et al.*, 2016; 2014a, 2004b, 2014c, 2013; Martelli *et al.*, 2016; Giampetruzzii *et al.*, 2016, 2015; Abbot, 2015; Fichtner *et al.*, 2015; Yaseen *et al.*, 2015; Alison, 2015; Simpson, 2015; Almeida and Nunney, 2015; EFSA, 2015; Djelouah *et al.*, 2014; Elbeaino *et al.*, 2014a, 2014b; White *et al.*, 2014; Frisullo *et al.*, 2014; Cariddi *et al.*, 2014; Boscia, 2014; Carlucci *et al.*, 2013); **California** (Krugner *et al.*, 2014, 2010; Krugner, 2010; Hernandez-Martinez *et al.*, 2007, Chen *et al.*, 2005; Costa *et al.*, 2004; Wong *et al.*, 2004); **Argentina** (Haelterman *et al.*, 2015) ; **Brazil** (Coletta-Filho *et al.*, 2016); **Florida** (Hopkins, 2014); **Spain** (Lopes *et al.*, 2014) ; **Portugal** (Pereira, 2015); **Slovenia** (Orešek and Seljak, 2015).

### **II-3 : Phytoplasma :**

**Italy** (Martelli *et al.*, 2013; 2000; 1999; Albanese *et al.*, 2012; Saponari and Savino, 2003; Pasquini *et al.*, 2000; Marzachi and Poggi Pollini, 2000; Marzachi *et al.*, 2000; Serrone *et al.*, 1996; Poggi Pollini *et al.*, 1996; Danielli *et al.*, 1996); **Albania** (Luigi *et al.*, 2009); **Iran** (Ghayeb Zamharir and Razavi, 2016; Ahangaran *et al.*, 2006); **Spain** (Font *et al.*, 1998) ; **Australia** (Spooner-Hart *et al.*, 2007).

### **III. RESULTS AND DISCUSSION**

Olive, like other crops, is not free from virological problems. The first report on a probable viral disease of olive goes back to 1938 (Pesante, 1938) and, since then, several virus-like diseases and viruses have been reported. Basing on all the available data in table 1, 15 viruses were reported in the olive tree. The most widespread were respectively Cucumber mosaic virus (CMV) and Olive leaf yellowing associated virus (OLYaV) in 19 countries for each; Followed by Olive latent virus-1 (OLV-1) (13 countries), Strawberry latent ring spot virus (SLRSV) and Arabis mosaic virus (ArMV) and Olive latent virus-2

(OLV-2) with 11 countries for each, Cherry leaf roll virus (CLRV) (10 countries), Olive latent virus-3 (8 countries), Olive latent ringspot virus (OLRSV) (8 countries), Tobacco necrosis virus D (TNV) (7 countries), Olive mild mosaic virus (OMMV) (5 countries), Olive vein yellows associated virus (OVYaV) (4 countries). Olive yellow mottling and decline associated virus (OYMDaV) (4 countries) and the less widespread viruses were Tobacco

mosaic virus (TMV) and Olive semil latent virus (OSLV) found in 3 countries for each. Martelli *et al.* (2002) have identified 14 virus species of 8 genera in olive trees, one more virus was not cited in their work, and it is OLV-3. This Virus was widespread in Syria, Turkey, Italy, Lebanon, Greece, Malta, Portugal, Syria and Tunisia.

Table.1: Geographical distribution of the olive tree viruses per continent:

	Europe	Africa	America	Asia	Australia – New Zealand
Olive leaf yellowing associated virus (OLYaV)	++++++ Spain-Italy-Croatia-Albania-Turkey-Greece-Cyprus-France	+++ Tunisia-Morocco-Algeria-Egypt	++ California-Chile	+++ Syria-Lebanon-Palestine-Israel	++ Australia
Cherry leaf roll virus (CLRV)	++++ Croatia- Italy- Spain-Portugal- Albania- Turkey	++ Egypt- Tunisia	----	++ Lebanon-Syria-	--
Strawberry latent ring spot virus (SLRSV)	++++ Croatia- Italy- Spain-Portugal- Turkey-Albania	++ Egypt- Tunisia	++ California	++ Lebanon-Syria	--
Arabis mosaic virus (ArMV)	++++++ Croatia- Italy- Spain-Portugal- Turkey- Croatia- Albania	++ Egypt Tunisia-	++ California	++ Syria-Lebanon-	--
Olive latent virus-1 (OLV-1)	++++++ Italy- Portugal- Turkey- Spain- Croatia- Albania	++ Egypt- Tunisia	++ California	++++ Jordan-Lebanon-Syria-Japan	--
Cucumber mosaic virus (CMV)	+++++++++ Croatia- Italy- Spain-Portugal- Albania-Turkey- France- Cyprus-Greece	+++ Egypt - Algeria-Tunisia Morocco	++ California	++++ Syria-Israel-Palastine Lebanon	++ Australia
Olive latent virus-2 (OLV-2)	++++++ Croatia- Italy- Greece-Portugal- Spain- Albania- Turkey-	++ Tunisia- Egypt	----	++ Syria-Lebanon	--
Olive latent virus-3 (OLV-3)	++++ Turkey- Italy- Greece-Malta- Portugal-	++ Tunisia	----	++ Syria-Lebanon	--
Olive semil latent virus (OSLV)	+++ Italy- Turkey- Portugal	----	----	---	---
Olive mild mosaic virus (OMMV)	++++ Portugal- Italy- Greece-Turkey- Spain	----	----	---	---

Olive yellow mottling and decline associated virus (OYMDaV)	++++ Portugal- Italy- Turkey- Spain	---	---	---	---
Olive latent ringspot virus (OLRSV)	++++ Croatia- Italy- Portugal- Turkey	++ Tunisia-Egypt	---	++ Syria- Lebanon	---
Olive vein yellows associated virus (OVYaV)	++++ Croatia- Italy- Portugal- Turkey	----	---	----	---
Tobacco necrosis virus D (TNV)	++++++ Croatia- Portugal- Italy- Albania- Turkey- Spain	----	+	----- California	----
Tobacco mosaic virus D (TMV)	+ Italy	----	+	+	Jordan ----

(+): Present;

(-): Absent

Our results were similar to those of Martelli (2013) that viruses of the olive tree were reported in 22 countries in five continents. All viruses were reported in Italy, some of these viruses were found only in the European continent, such as Olive vein yellows associated virus (OVYaV), Olive yellow mottling and decline associated virus (OYMDaV), Olive mild mosaic virus (OMMV) and Olive semilatent virus (OSLV) (Table 1). Cucumber mosaic virus (CMV) and Olive leaf yellowing associated virus (OLYaV) were in the five continents (Table 1), these two viruses were the only ones reported to the olive tree in Australia (Table 1). We can explain this by the far of this continent to the others that may lead to decrease the dissemination vectors and factors, or it may caused by the virulence adaptation of these viruses on the olive trees in Australia.

Virus Olive trees are infected with systemic pathogens such as viruses and phytoplasmas as well as agents of diseases of unknown etiology (Çağlayan *et al.*, 2011). The first report of a probable viral disease of olive (*Olea europaea* L.) trees goes back to 1938 (Pesante, 1938). Since then, several viruses and phytoplasmas were reported in the Mediterranean countries where olive is economically important and olive oil is considered one the main components of the Mediterranean diet (Çağlayan *et al.*, 2011).

Several olive tree diseases caused by viruses were reported from several countries, viruses were mechanically transmitted to indicator plants and of the seven that were subsequently identified, and four belong to the nepovirus group: Strawberry latent ringspot (SLRV), cherry leaf roll (CLRV), arabis mosaic (AraMV) and olive latent ringspot (OLRV); one belongs to the cucumovirus group: cucumber mosaic virus 'CuMV' and two others have not been assigned to any of the existing groups: olive latent virus-1 (OLV-1) and olive latent virus-2 (OLV-2). Sometimes virus are detected in *Olea*

plants that show symptoms, as severely distorted leaves and misshapen fruits – which are particularly evident in infected olive trees in Portugal and in Italy (da clara henriques, 1994)

SLRV was found to be associated to a severe symptomatology of Italian cv. Ascolana tenera, consisting of distorted sickle shaped leaves and fruit deformation (Marte *et al.*, 1986). Leaf symptoms were transmitted to healthy Ascolana plants and virus recovered from these. This work clearly showed a correlation between field symptoms and SLRV infection. Identical severe leaf and fruit symptoms were observed in a Portuguese table olive cv. Negrinha, where SLRV was also diagnosed (Henriques *et al.*, 1990; 1992). In Addition, this virus was also detected in olive plants previously shown to contain virus-like particles in pollen, as well as in apparently normal plants (Savino *et al.*, 1979; Henriques *et al.*, 1992).

ArMV was isolated from declining trees with sickle shaped leaves (Savino *et al.*, 1979) and OLRV from plants showing partial defoliation (Savino *et al.*, 1983). CLRV and CMV were recovered from symptomless trees (Savino and Gallitelli, 1983; Rei *et al.*, 1993). OLV-1 was isolated from an olive tree showing some fasciations and bifurcation of leaves and stems (da Clara Henriques, 1994).

OLV-2 was recovered from symptomless olive trees. The virus particles possess a distinctive bacilliform shape which readily differentiates it from known viruses (Martelli and Gallitelli, 1985). Absence of symptoms in olive trees is by no means a reliable indication of absence of infection. As mentioned before, almost all olive virus isolates could be recovered from assymptomatic trees, with the exception of SLRV infecting Ascolana tenera and Negrinha cvs., where the virus is associated to severe symptoms.

The epidemiology of most olive tree viruses is still unknown. Although some of these viruses are soil-borne (SLRSV, ArMV, TNV) others can be transmitted mechanically (TMV), by seed (CLRV and OLV-1) (Saponari *et al.*, 2002a), by aphids (CMV) or only by mechanical inoculation and grafting (OLV-2 and OLRSV). However, there is very limited knowledge about virus transmission mechanism(s) under field conditions. OLYaV was reported to be detected in mealybugs and psyllids that fed on an infected olive tree (Sabanadzovic *et al.*, 1999) but it is too early to regard these insects as vectors of the virus because preliminary transmission trials were negative.

The fact that olive viruses occur in latent forms, makes it impossible to visually assess infected trees and determine their impact on plant yield. Nepoviruses are known to infect important crops, including grapevine, often grown

in close proximity to olives in Mediterranean countries, where they cause considerable losses (da clara henriques, 1994). Cucumoviruses have a wide range of hosts, some of high economic importance and may cause severe crop damage (da Clara Henriques, 1994). Thus, it is plausible to respect that viruses of these groups cause deleterious effects on olive trees (da clara henriques, 1994). Severe fruit deformations are associated to SLRV infections in table olive cvs., thus leading to a commercial devaluation of the product. SLRV is also associated to a poor rooting ability of the olive cuttings (Henriques *et al.*, 1992). Olive viruses may also negatively affect vigour, longevity and ability of olive trees to stand environmental stress (da clara henriques, 1994).

Table.2: Geographical distribution of the pathogenic bacteria reported to the olive tree per continent:

	Europe	Africa	America	Asia	Australia – New Zealand
<i>Pseudomonas savastanoi</i> pv. <i>savastanoi</i>	++++++ Spain- Italy- France- Portugal- Greece- Turkey Croatia	++++ Tunisia- Algeria- Morocoo- Egypt	+\nCalifornia	++++++ Jordan- Lebanon- Iraq- Nepal- Iran- Syria- China- Japan	++\nNew Zealand- Australia
<i>Xylella fastidiosa</i>	++++ Italy- Spain- Portugal- Slovenia	----	+++ USA- Argentina- Brazil	----	----

(+): Present;

Olive knot, caused by *Pseudomonas savastanoi* pv. *savastanoi* (Pss) is a serious disease of olive (*Olea europaea* L.) in the Mediterranean (Krid *et al.*, 2010). Basing in the available data, this bacteria was detected in 21 countries in five continents, the majority of these countries exist in the Mediterranean Basin. The disease characteristic symptoms are knots that develop as a reaction to factors produced by the bacterium, including indole-3-acetic acid (IAA) (Matas *et al.*, 2009; Rodríguez-Moreno *et al.*, 2008), cytokinins (Surico *et al.*, 1985), products of the type III secretion system (TTSS), encoded by the hrp/hrc gene clusters (Sisto *et al.*, 2004) and a variety of genes encoding known *P. syringae* virulence determinants (Pérez-Martínez *et al.*, 2008). Olive knot disease is characterized by hyperplasia formation on the stems and branches of olive plants and occasionally on the leaves and fruits (Surico, 1986). The disease can cause severe damage in olive groves, mainly when weather conditions favor the survival of epiphytic populations of the pathogen and their entry into the bark. *P. syringae* subsp. *savastanoi* has epiphytic resident

(-): Absent

populations on olive twigs, leaves, and drupes, with larger populations on twigs than on leaves as a result of the better survival of the pathogen in the rough bark surface (Lavermicocca, and Surico. 1987).

*X. fastidiosa* subsp. was detected in Europe in four countries (Italy, Spain, Portugal and Slovenia) and in America in three countries (USA, Argentina and Brazil) (Table 2). It is endemic to North America only and infects numerous hosts, but generally not grapevine (Davis *et al.*, 1978; Hopkins and Mollenhauer, 1973). *X. fastidiosa* subsp. *pauca* contains South American strains causing citrus variegated chlorosis and coffee leaf scorch (Hopkins and Purcell, 2002). *X. fastidiosa* subsp. *sandyi* comprises closely related strains isolated from oleander in California and Texas (USA). As to olive (*Olea europaea*), there are reports from California, where the bacterium was detected and/or isolated from trees with leaf scorching (Wong *et al.*, 2004), unspecified symptomatology, if any (Hernandez-Martinez *et al.*, 2007), or showing dieback and leaf scorch (Krugner *et al.*, 2010). Its properties, biology, epidemiology and disease

management strategies have been exhaustively described in a number of reviews to which the readers are referred for more detailed information (Purcell, 2013; Janse and Obradovic, 2010; Chatterjee *et al.* 2008; Hopkins 1989; Purcell and Hopkins 1996; Redak *et al.* 2004).

The quick decline syndrome of olive (OQDS) caused by *Xylella fastidiosa* is a destructive disorder that appeared suddenly a few years ago in the groves of a restricted area of the Apulian province of Lecce, Salento peninsula, south-east Italy (Martelli, 2013). The disease appeared suddenly in the autumn and spring, mainly on the lower branches of the trees and was particularly severe on "Ogliarola di Lecce", the olive cultivar most widely grown in the area, whereas "Cellina di Nardò", the other major local cultivar, appeared to be resistant (Frisullo *et al.*, 2014), and in Argetina olive trees older than 50 years of age were observed in six orchards of cv. Arauco, which showed symptoms recalling those induced by *Xylella fastidiosa*. Some branches displayed desiccated leaves at the top and basal leaves with apical scorching. Additional symptoms were slow decay, dull green coloration, curling and necrosis of the leaves, partial defoliation and rapid death of shoots and branches.

On this basis, the phytoplasmas identified in olive trees have been classified as 16S-IB (Aster yellows group), 16S-VA (Elm yellows group) and 16S-XIIA (Stolbur group) (Pasquini *et al.*, 2000). After the first report in Italy in 1995, several reports confirmed the association between olive foliage disorders and the presence of phytoplasmas in Spain, Italy and Iran (Font *et al.*, 1998; Pasquini *et al.*, 2000; Ahangaran *et al.*, 2006).

Phytoplasma was detected on the olive tree in five countries, it was responsible of yellow symptoms in a few branches was observed. Some leaves showed diffuse chlorotic mottling affecting almost all the blade, while others showed mottling with green and yellow areas: leaf deformation was also sometimes present (Poggi Pollini *et al.*, 1996). Affected trees exhibited a rather variable range of symptoms: shoot proliferation, leaf rolling, yellowing of leaves and branches, phyllody, fasciation, witches'-brooms, dwarfing, leaf bronzing, abortion of axillary buds, erect growth, appearance of small heads due to clustering of hypertrophoyd inflorescences, decline and die-back (Pasquini *et al.*, 2000).

#### IV.CONCLUSION

Olive tree is an important crop, but it is unfortunately attacked by several pathogens, Fungi, bacteria, viruses, phytoplasmas...., these diseases are widespread in the five continent. 15 viruses were cited as pathogens of the olive tree, showing several symptoms. Some of the existed in all continents on different olive tree varieties, others they were reported just in few countries in the same region or

same continent or even in just one country. Two bacteria were cited on the olive tree, which show big damage to the tree such as knots, dieback, decline, defoliation and chlorosis. After this work, we should have a prevention control with diagnostics of to the olive trees coming from other countries caring these pathogens to another country and prevention treatments against bacteria.

#### REFERENCES

- Abbott A., 2015. Italian scientists vilified in wake of olive-tree deaths. *Nature* 522, 13–14.
- Abdine M., Allak H., Nus B.E., Nigro F., Catalano L., Digiaro M. Phytosanitary aspects and nursery production of olive in Syria: pitfalls and perspectives [printed version also available in Arabic]. In: Di Terlizzi B. (ed.), Dragotta A. (ed.), Jamal M. (ed.). Syrian national strategic plan for olive oil quality: final report. Bari: CIHEAM, 2007. p. 143-151.
- Abou Ghanem N., Sabanadzovic M., Mahboubi B., Terlizzi Di, Sabanadzovic S., Savino V., Uyemoto J.K., Martelli G.P., 2001. Molecular detection of a closterovirus associated with Apricot stem pitting in southern Italy. *Journal of Plant Pathology*. 83 (2): 125-132.
- Abu-Ghorrah, M. (1988). Taxonomie et pouvoir pathogène de *Pseudomonas syringae* pv. *savastanoi*. Thesis (Ph. D.). University of Angers, France
- Adam DB, Pugsley AT (1934) Bacterial diseases in Victoria. *Journal of Agriculture, Victoria*, 32: 304-311. 323.
- Adaskaveg J.E., Forster H., Thompson D., Nguyen K., Connel J., 2012. Epidemiology and management of olive knot caused by *Pseudomonas syringae* pv. *savastanoi*. Annual project/research grant report 66-75.
- Ahmad A.A., Moretti C., Valentini F., Hosni T., Farag N.S., Galal A.A., Buonauro R., 2009. Olive knot caused by *Pseudomonas savastanoi* pv. *savastanoi* in Egypt. *Journal of Plant Pathology*. 91 (1), 231-240.
- Al Abdullah A, Minafra A, Elbeaino T, Saponari M, De Stradis A, Savino V, Martelli GP. (2010) Complete nucleotide sequence of Olive latent virus 3, representing a new genome structure in the genus Marafivirus. *Virus Res* 152:10–17.
- Al Abdullah A., 2004. Preliminary evaluation of the sanitary status of olive in Syria. <http://agris.fao.org/agris-search/search.do?recordID=QC2004002094>
- Al Rwahnih M., Guo Y., Daubert S., Golino D. and Rowhani A., 2011. Characterization of latent viral infection of olive trees in the national clonal germplasm repository in California. *Journal of Plant Pathology*. 93 (1), 227-231.
- Gómez-Lama C.C., Sesmero R., Valverde-Corredor A., Javier López-Escudero F., Mercado-Blanco J., 2017. A

- split-root system to assess biocontrol effectiveness and defense-related genetic responses in above-ground tissues during the tripartite interaction *Verticillium dahliae*-olive-*Pseudomonas fluorescens* PICF7 in roots. *Plant and Soil.* 1–20.
- Alabdalla N., Valentini F., Moretti C., Essa S., Buonauro R., Abu-Ghorra M., 2009. First report of *Pseudomonas savastanoi* pv. *savastanoi* causing olive knot in Syria. *Plant pathology.* 58(6): 1170.
- Al-abdallah O., Elbeaino T., Farinelli D., Valentini F., Digiaro M., 2010. Sanitary assessment of olive varieties in a collection plot at the University of Perugia (Italy). 2<sup>nd</sup> Conference on Biotechnology Research and Applications in Palestine, 26-27th September, Page 34
- Albanese G., Faggioli F., Ferretti L., Sciarroni R., La Rosa R., Barba M., 2003. Sanitary status evaluation of olive cultivars in Calabria and Sicily. *Journal of Plant Pathology* 85: 304.
- Alison A., 2015. Scientists blamed for olive-tree ruin: Italian police investigate researchers' role in a bacterial epidemic that is devastating Puglia's olive groves. *Nature*, 522 (7544): 13.
- Alkowni R (2000) Developing and application of molecular diagnosis for the detection of olive-infecting viruses and molecular characterization of *Olive latent ring spot virus* (OLRSV). PhD Thesis, University of Bari (IT).
- Alkowni R, Grieco F., Martelli GP (2000a) Complete nucleotide sequence of RNA-2 of olive latent ringspot virus. *Archives of Virology.* 145: 1-7.
- Almeida R.P.P., 2016. Can Apulia's olive trees be saved? *Science*, 353 (6297): 346-348. DOI: 10.1126/science.aaf9710
- Almeida, R.P.P. and Nunney, L. (2015). How do plant diseases caused by *Xylella fastidiosa* emerge? *Plant Disease*, 99, 1457-1467.
- Alvarez F, Garcia de los Rios JE, Jimenez P, Rojas A, Riche P, Troya MT, 1998. Phenotypic variability in different strains of *Pseudomonas syringae* subsp. *savastanoi* isolated from different hosts. *European Journal of Plant Pathology* 104, 603–9.
- Aragon I.M., Perez-Martinez I., Moreno-Perez A., Cerezo M. Ramos C., 2014a. New insights into the role of indole-3-acetic acid in the virulence of *Pseudomonas savastanoi* pv. *Savastanoi*. *FEMS Microbiol Lett*, 356: 184–192.
- Aragón I.M., Pérez-mendoza D., Gallegos M.T., and Ramos C., 2014b. The c-di-GMP phosphodiesterase BifA is involved in the virulence of bacteria from the *Pseudomonas syringae* complex. *Molecular plant pathology.* 1-12.
- Aragón, I.M., Pérez-Mendoza, D., Gallegos, M.T., and Ramos, C. (2015). The c-di-GMP phosphodiesterase BifA is involved in the virulence of bacteria from the *Pseudomonas syringae* complex. *Mol.PlantPathol.* doi:10.1111/mpp. 12218.
- Ark, P.A., Thompson, J.P., 1960. Experimental greenhouse control of crown gall and olive knot with antibiotic drenches. - *Plant Dis. Repr.* 44: 197-199.
- Azeri T., 1993. Research on olive leaf spot, olive knot and verticillium wilt of olive in Turkey. detin OEIP/EPPO Bulletin 23,437-440.
- Balestra GM, Varvaro L (1997) Influence of nitrogen fertilization on the colonization of olive phylloplane by *Pseudomonas syringae* subsp. *savastanoi*. In: Rudolph K (ed) *Pseudomonas syringae* pathovars and related pathogens. Kluwer Academic Publishers, Dordrecht, pp 88–92.
- Balestraa G.M., Lamichhanea J. R., Kshetrib M. B., Mazzagliaa A., Varvaro L., 2009. First report of olive knot caused by *Pseudomonas savastanoi* pv. *savastanoi* in Nepal. *Plant Pathology*, 58, 393.
- Baratta B, Di Marco L (1981) Control of olive knot attacks on the cultivar Nocellara del Belice. *Informatore Fitopatologico* 31, 115–116.
- Barba M., 1993. Viruses and virus-like diseases of olive. *Bulletin OEPP/EPPO Bulletin* 23, 493-497.
- Bardaji, L., Pérez-Martínez, I., Rodríguez-Moreno, L., Rodríguez-Palenzuela, P., Sundin, G. W., Ramos, C., et al. (2011). Sequence and role in virulence of the three plasmid complement of the model tumor-inducing bacterium *Pseudomonas savastanoi* pv. *savastanoi* NCPPB 3335. *PLoS ONE* 6:e25705.
- Bartolini, S., Saponari, M., Guerriero, R. and Loreti, F. (2002). Two new clones of cultivar 'Leccino'. *Acta Hortic.* 586, 225-228 DOI: 10.17660/ActaHortic.2002.586.42
- Basim E., Basim H., Ozcan M., 2006. Antibacterial activities of Turkish pollen and propolis extracts against plant bacterial pathogens. *Journal of Food Engineering* 77: 992–996.
- Basim H., Ersoy A., 2001. Identification of *Pseudomonas savastanoi* pv. *savastanoi*, olive knot pathogen, by polymerase chain reaction. *Phytopathology* 91: S 6.
- Basim, H., 2002. Characterization by Pulsed-field Gel Electrophoresis of *Pseudomonas savastanoi* pv. *savastanoi* strains in Western Mediterranean Region of Turkey. APS. SO. MSA Joint Meeting, August 25- 29 Salt Lake, USA. 6.
- Basim, H.O. Yeğen, Ersoy, Ülger., 2000. Batı Akdeniz Bölgesinde Zeytin Ağacı'nda *Pseudomonas savastanoi* pv. *savastanoi* Tarafından Oluşturulan Bakteriyel Dal Kanseri Hastalığının Yayılışı ve Hastalık Etmeninin Tanısı. Türkiye 1. Zeytinçilik sempozumu 6-9 Haziran 2000 Uludağ üniversitesi Ziraat Fak. Bahçe Bit. ve Gıda Mühendisliği Bölümleri, Bursa, 310-315s.

- Beler Ö., Açıkgöz S., 2005. Determining of Some Virus Diseases Using the ELISA Test in Olive Nurseries and Olive Trees in the Region of Aegean and Marmara. ADÜ Ziraat Fakültesi Dergisi 2005; 2(1): 79 – 84.
- Bellini, E., Giordani, E. and Parlati, M.V. (2002). Three new olive cultivars obtained by cross-breeding. Acta Hortic. 586, 221-223. DOI: 10.17660/ActaHortic.2002.586.41
- Beltra, R. 1958. Relacion entre la concentración de triptofano en los tallos do olivo y la localizacion de los tumores bacterianos. Microbiol. Espan. 11:401-410. 2.
- Beltra, R. 1961. Efecto morfogenetico observado en los extractos hormonales de los tumores del olivo. Microbiol. Espan. 14:177-187.
- Benjama A, Walali L, Jananti L, Moukhli A (1987) Field reaction of different varieties of olive (*Olea europaea* L.) to olive knot disease caused by *Pseudomonas syringae* pv. *savastanoi* (E.F. Smith) Stevens. Third International Working Group on *Pseudomonas syringae* pathovars. Lisbon, Portugal, 1–4 September 1987. pp. 41–52.
- Benjama A., 2003. Méthode d'évaluation rapide du degré d'attaque de l'olivier par la tuberculose cause par *Pseudomonas savastanoi* pv. *savastanoi* en verger au Maroc. Fruits, 58, 213-219.
- Benjama A., Atbi M., Tahiri J., Aznadi M., Situation actuelle de la tuberculose de l'olivier dans la région de Zerhoun au Maroc, El- Awamia 83 (1993) 153–160.
- Benjama A., Boubaker A., Khlef H., Krimi Z., Nesme X., Lopez M., Zoina A., Susceptibility of some stone fruit rootstocks and rose to crown-gall, caused by *Agrobacterium tumefaciens* in the Arab Mediterranean countries, B. OEPP 32 (3) (2002) 463–466.
- Benjama A., Gardan L., Influence du milieu de culture et du temps d'incubation sur la production d'AIA par *Pseudomonas savastanoi* pv. *savastanoi*, El-Awamia 88 (1995) 79–92.
- Benjama A., Isolement et identification de l'agent pathogène de la tuberculose de l'olivier au Maroc, El-Awamia 72 (1989) 57–69.
- Benjama A., La tuberculose de l'olivier au Maroc, Fiche technique DIF – Inra, Maroc, 2002.
- Benjama A., Parasitic olive diseases in Morocco, Olivae 20 (1988) 21–33.
- Benjama A., Preliminary studies on olive knot in Morocco caused by *Pseudomonas syringae* pv. *savastanoi*., Arab. J. Plant Protect. 5 (2) (1987) 67–69.
- Benjama A., Walali L., Janati L., Moukhli A., Field reaction of different varieties of olive *Olea europaea* L. to olive knot disease caused by *Pseudomonas syringae* pv. *Savastanoi* (E.F. Smith) Stevens, El-Awamia 75 (1992) 41–52.
- Benjama, A. 1994. Étude de la sensibilité variétale de l'olivier au Maroc vis-à-vis de *Pseudomonas syringae* pv. *savastanoi*, agent de la tuberculose. Cahiers Agric. 3:405-408.
- Benjama, A., 1990. Isolement et identification de l'agent pathogène de la tuberculose de l'olivier au Maroc. Al Awamia, 72 : 57-69.
- Bertolini E, Olmos A, Martínez MC, Gorris MT, Cambra M (2001a) Single-step multiplex RT-PCR for simultaneous and colourimetric detection of six RNA viruses in olive trees. J Virol Methods 96: 33–41.
- Bertolini E, Penyalver R, Garcia A et al., 2003. Highly sensitive detection of *Pseudomonas savastanoi* pv. *savastanoi* in asymptomatic olive plants by nested-PCR in a single closed tube. Journal of Microbiological Methods 52, 261–6.
- Bertolini E., Cambra M., Peñalver R., Ferrer A., Garcia A., Del Rio M.C., Gorris M.T., Martínez M.C., Quesada J.M., García De Oteyza J., Duran-Vila N., Caballero J.L., López M.M., 2001b. Métodos serológicos y moleculares de diagnóstico de virus y bacterias de olivo. Evaluación de la sensibilidad varietal y la aplicación a programas de certificación. Mercacei Magazine 30:1-6.
- Bertolini, E. (2003). Virosis y bacteriosis del olivo: detección serológica y molecular. Thesis (Ph. D.). Universidad Politécnica de Valencia, Spain.
- Bertolini, E., Olmos, A., Lopez, M.M., & Cambra, M. (2003b). Multiplex nested reverse transcription-polymerase chain reaction in a single tube for sensitive and simultaneous detection of four RNA viruses and *Pseudomonas savastanoi* pv. *savastanoi* in olive trees. Phytopathology, Vol. 93, No. 3, (March 2003), pp. 286-292, ISSN 0031-949X.
- Bertolini, E., Penyalver, R., García, A., Olmos, A., Quesada, J.M., Cambra, M., & López, M.M. (2003a). Highly sensitive detection of *Pseudomonas savastanoi* pv. *savastanoi* in asymptomatic olive plants by nested-PCR in a single closed tube. Journal of Microbiological Methods, Vol. 52, No. 2, (February 2003), pp. 261-266, ISSN 0167-7012.
- Bertolini, E.; Fadda, Z.; Garcia, F.; Celada, B. ; Olmos, A. ; Gorris, M. T. ; Del Rio, C. ; Caballero, J. ; Duran-Vila, N. ; Cambra, M. ; 1998 : Virosis del olivo detectadas en Espana. Nuevos metodoa de diagnostico. Phytoma Espana 102, 191-193.
- Moretti C., Vinatzer B.A., Onofri A., Valentini F., Buonauro R., 2017. Genetic and phenotypic diversity of Mediterranean populations of the olive knot pathogen, *Pseudomonas savastanoi* pv. *savastanoi*. 66 (4): 595–605.
- Bianco A., Studies in Natural Products Chemistry, vol. 32, Elsevier Science, Amsterdam, The Netherlands, 1990.
- Bjeliš, M., Loconsole G. i Saponari M., 2007. Nazočnost virusa u maslinicima Hrvatske. Pomologia Croatia. 13 (3): 165-172.

- Bleve G., Marchi G., Ranaldi F., Gallo A., F. Cimaglia, A. F. Logrieco, G. Mita, Ristori J., and Suri G., 2016. Phytopathologia Mediterranea. Molecular characteristics of a strain (Salento-1) of *Xylella fastidiosa* isolated in Apulia (Italy) from an olive plant with the quick decline syndrome. 55: 1: 139–146.
- Boscia D., 2014. Occurrence of *Xylella fastidiosa* in Apulia. Journal of Plant Pathology (2014), 96 (4): 97-98.
- Bosso L., Russo D., DI Febbraro M., Cristinzio G. and Zo A., 2016. Potential distribution of *Xylella fastidiosa* in Italy: a maximum entropy model. Phytopathologia Mediterranea. 55 (1): 62–72.
- Botalico, A., and Ercolani, G. L. 1971. *Pseudomonas savastanoi* (E.F. Smith) Stevens su Ligusto giapponese in Puglia. Phytopathol. Mediterr. | (): 132-135. 4.
- Bottalico G., Rodio M.E., Saponari M., Savino V. and Martelli G.P. (2002). Preliminary results of sanitation trials of viruses infecting olive. Journal of Plant Pathology. 84, 176.
- Bouaichi A., Benkirane R., Ameur A., Sadik S., Benbouazza A., and Achbani E.H., 2015. Characterization of *Pantoea agglomerans* Strains Isolated from Olive Knot disease in Morocco. Biolife. 3(2):533-539.
- Boulila M., 2016. Investigations of molecular evolutionary mechanisms in partially sequenced heat shock protein70 homologue-coding gene of Olive leaf yellowing-associated virus isolates from Tunisia. Journal of DNA and RNA Research. 1 (1): 1-30.
- Boulila M., Mahjoub M., 1994. Inventaire des maladies de l'olivier en Tunisie. EPPO Bulletin. 24 (4) : 817–823.
- Boulila, M., Mahjoub, M., Romdhani, M.S. & Ben Othman, M.N. (1993) La maladie du pourridié-agaric observée dans des oliveraies tunisiennes. Bulletin OEPP/EPPO Bulletin 23, 447–448.
- Bozkurt I.A., Soylu S., Mirik M., Ulubas Serce C. and Baysal O., 2014. Characterization of bacterial knot disease caused by *Pseudomonas savastanoi* pv. *savastanoi* on pomegranate (*Punica granatum* L.) trees: a new host of the pathogen. Letters in Applied Microbiology 59, 520—527.
- Bracci T., M. Busconi, C. Fogher, and L. Sebastiani, “Molecular studies in olive (*Olea europaea* L.): overview on DNA markers applications and recent advances in genome analysis,” Plant Cell Reports, vol. 30, no. 4, pp. 449–462, 2011.
- Braithwaite M, Alexander BJR, Young JM, Ganev S (1999) Olive knot disease in New Zealand. In ‘Proceedings of the 12<sup>th</sup> Australasian Plant Pathology Society Conference’. Canberra, 1999. (Ed. L. Morin) p. 157.
- Brown, N. A. 1932. Canker of ash trees produced by a variety of the olive tubercle organism, *Bacterium savastanoi*. J. Agric. Res. 44:701-722.
- Buonauro R., Moretti C., Passos da Silva D., Cortese C., Ramos C. and Venturi V., 2015. The olive knot disease as a model to study the role of interspecies bacterial communities in plant disease. Frontiers in Plant Science. 6 :1-12.
- Çağlayan K., Faggioli F., Barba M., 2011. Viruses, Phytoplasmas and Diseases of Unknown Etiology of Olive Trees. In book: Virus and virus-like diseases of pome and stone fruits, Chapter: 53, Publisher: The American Phytopatological Society, Editors: A. Hadidi, M. Barba, T. Candresse, W. Jelkmann, pp.289–297.
- Çağlayan K., Fidan U., Tarla G., and Gazel M. 2004, First report of olive viruses in Turkey. J. Plant Pathol., 86 (1): 89-90.
- Campos A., da Costa G., Varela Coelho A., Fevereiro P., 2009. Identification of bacterial protein markers and enolase as a plant response protein in the infection of *Olea europaea* subsp. *europaea* by *Pseudomonas savastanoi* pv. *Savastanoi*. Eur. J. Plant Pathol., 125:603–616.
- Capasso, R., Cristinzio, G., Evidente, A., Visca, C., and Iannini, C. (1997). “Oleuropein and other polyphenols from olive (*Olea europaea*) for protecting the plant against *Pseudomonas syringae* subsp. *savastanoi*,” in *Pseudomonas syringae* Pathovars and Related Pathogens, eds K. Rudolph, T. J. Burr, J. W. Mansfield, D. Stead, A. Vivian, and J. von Kietz (Dordrecht: Kluwer Academic Publishers), 133–137.
- Capasso, R., Evidente, A., Schivo, L., Orru, G., Marcialis, M. A., and Cristinzio, G. (1995). Antibacterial polyphenols from olive oil mill waste-waters. J. Appl. Bacteriol. 79, 393–398.
- Caponero A, Contesini AM & Iacobellis NS (1995) Population diversity of *Pseudomonas syringae* subsp. *savastanoi* on olive and oleander. Plant Pathol 44: 848–855.
- Caballo-Ponce E., Van Dillewijn P., Wittich R.M., Ramos C., 2017. WHOP, a Genomic Region Associated with Woody Hosts in the *Pseudomonas syringae* Complex Contributes to the Virulence and Fitness of *Pseudomonas savastanoi* pv. *savastanoi* in Olive Plants. International Society for Molecular Plant-Microbe Interactions. 30 (2): 113-126.
- Capuzzo C., Firrao G., Mazzon L., Squartini A. and Girolami V., 2005. ‘Candidatus *Erwinia dacicola*’, a coevolved symbiotic bacterium of the olive fly *Bactrocera oleae* (Gmelin). International Journal of Systematic and Evolutionary Microbiology, 55 : 1641–1647.
- Cardoso J.M.S., Félix M.R., CLARA M.I.E., Oliveira S., 2012. First characterization of infectious cDNA clones of Olive mild mosaic virus. Phytopathologia Mediterranea. 51 (2): 259–265.

- Cardoso J.M.S., Félix M.R., Clara M.I.E., Oliveira S., 2012. First characterization of infectious cDNA clones of Olive mild mosaic virus. *Phytopathologia Mediterranea*. 51 (2): 259-265.
- Cardoso J.M.S., M.R. Felix, S. Oliveira and M.I.E. Clara, 2004. A tobacco necrosis virus D isolate from *Olea Europea* L.: viral characterization and coat protein sequence analysis. *Archives of Virology* 149, 1129–1138.
- Cardoso, J. M. S., Félix, M. R., Clara, M. I. E., & Oliveira, S. (2005). The complete genome sequence of a new necrovirus isolated from *Olea europaea* L. *Archives of Virology*, 150, 815–823.
- Cardoso, J. M. S., Félix, M. R., Clara, M. I. E., & Oliveira, S. (2009). Complete genome sequence of a Tobacco necrosis virus D isolate from olive trees. *Archives of Virology*, 154, 1169–1172.
- Cariddi C., Saponari M., Boscia D., De Stradis A., Loconsole G., Nigro F., Porcelli F., Potere O. and Martelli G.P., 2014. Isolation of a *Xylella fastidiosa* strain infecting olive and oleander in Apulia, Italy. *Journal of Plant Pathology*, 96 (2), 425-429.
- Carlucci A., Lops F., Marchi G., Mugnai L., Surico G., 2013. Has *Xylella fastidiosa* "chosen" olive trees to establish in the Mediterranean basin?. *Phytopathologia Mediterranea*. 52 (3): 541-544.
- Casano, F.J., 1985. Tentativi di identificazione sierologica di isolati di *Pseudomonas syringae* pv. *savastanoi*. Annali 1st. spcr. Patol. veg. Roma 10,23-28.
- Castellano M.A., Difranco A., Martelli G.P., 1987. Electron microscopy of two olive viruses in host tissues. *Journal of Submicroscopic cytology*. 19: 495-508.
- Castellano, M. A., Loconsole, G., Grieco, F., Di Sansebastiano, G. P., & Martelli, G. P. (2005). Subcellular localization and immunodetection of movement proteins of olive latent virus 1. *Archives of Virology*, 150, 1369–1381.
- Catara, V., Colina, P., Bella, P., Tessitori, M., & Tirrò, A. (2005). Variabilità di *Pseudomonas savastanoi* pv. *savastanoi* in un'area olivicola della Sicilia e comportamento di alcune varietà di olivo alle inoculazioni [*Olea europaea* L.]. *Tecnica Agricola*, Vol. 57, No. 1/2, (January-June 2005), pp. 41-52, ISSN 0371-5124.
- Cayuela, J. A., Rada, M., Rios, J. J., Albi, T., and Guinda, A. (2006). Changes in phenolic composition induced by *Pseudomonas savastanoi* pv. *savastanoi* infection in olive tree: presence of large amounts of verbascoside in nodules of tuberculosis disease. *J. Agr. Food Chem.* 54, 5363–5368.
- Çağfçg O., Serçe Ç.U., 2014. Research on the Presence of Some Viruses in Olive Orchards Using Serological and Molecular Techniques in Mardin Province of Turkey. *Bitki Koruma Bülteni* 2014, 54(3):233-246.
- Chatterjee S., Almeida R.P.P, Lindow S., 2008. Living in two Worlds: The plant and insect lifestyles of *Xylella fastidiosa*. *Annual Review of Phytopathology* 46: 243-271
- Chen, J., Groves, R., Civerolo, E. L., Viveros, M., Freeman, M., and Zhang, Y. 2005. Two *Xylella fastidiosa* genotypes associated with almond leaf scorch disease on the same location in California. *Phytopathology* 95:708-714.
- Caglayan K., Faggioli F., Barba M., 2011. Viruses, phytoplasmas and diseases of unknown etiology of olive trees. [https://www.researchgate.net/publication/241687505\\_Viruses\\_phytoplasmas\\_and\\_diseases\\_of\\_unknown\\_etiology\\_of\\_olive\\_trees](https://www.researchgate.net/publication/241687505_Viruses_phytoplasmas_and_diseases_of_unknown_etiology_of_olive_trees).
- Chliyeh M., Rhimini Y., Selmaoui K., Ouazzani Touhami A., Filali-Maltouf A., El Modafar C., Moukhli A., Oukabli A., Benkirane R., Douira A., 2014. Comparative study of pathogenicity tests for *Verticillium dahliae* and *Phytophthora palmivora* causing wilt and decline of olive tree (*Olea europaea* L.). *Int. J. Pure App. Biosci.* 2 (2): 28-38.
- Choueiri E., Freiji, A. Kubaa, R. A.; Elbeaino, T. 2015. First report of Cucumber mosaic virus and Olive latent ringspot virus on olive trees in Lebanon. *Journal of Plant Pathology*. 97 (1): 221.
- Ciccarone A., 1950. Alterazioni da freddo e da roagna sugli ulivi, esemplificate dai danni osservati in alcune zone pugliesi negli anni 1949-1950. *Bollettino della Stazione di Patologia Vegetale di Roma* 6: 141-174.
- Cimmino A., Marchi G., Surico G., Hanuszkiewicz A., Evidente A., Holst O., 2008. The structure of the O-specific polysaccharide of the lipopolysaccharide from *Pantoea agglomerans* strain FL1Carbohydrate Research Volume 343, Issue 2, 392–396
- Cimmino A., Andolfi A., Marchi G., Surico G., and Evidente A., 2006. Phytohormone production by strains of *Pantoea agglomerans* from knots on olive plants caused by *Pseudomonas savastanoi* pv. *Savastanoi*. *Phytopathol. Mediterr.* (2006) 45, 247–252.
- Cinelli T., Moscetti I., Marchi G., 2014. PsasM2I, a Type II Restriction–Modification System in *Pseudomonas savastanoi* pv. *savastanoi*: Differential Distribution of Carrier Strains in the Environment and the Evolutionary History of Homologous RM Systems in the *Pseudomonas syringae* Complex. *Microb Ecol.*, 68:842–858.
- Cinelli T., Rizzo D., Marchi G. Surico G., 2013. First Report of Knot Disease Caused by *Pseudomonas savastanoi* on Sweet Olive in Central Italy. *Disease Notes*. 97 (3): 419.
- Coletta-Filho H.D., Francisco C.S., Lopes J.R.S., De Oliveira A.F., Da Silva L.F.D.O., 2016. First report of olive leaf scorch in Brazil, associated with *Xylella*

- fastidiosa* subsp. *Pauca*, *Phytopathologia Mediterranea*. 55(1): 130–135.
- Comai L & Kosuge T (1980) Involvement of plasmid deoxyribonucleic acid in indoleacetic acid synthesis in *Pseudomonas savastanoi*. *J Bacteriol* 143: 950–957.
- Comai L., G. Surico, Kosuge T., 1982. Relation of Plasmid DNA to Indoleacetic Acid Production in Different Strains of *Pseudomonas syringae* pv. *Savastanoi*. *Journal of General Microbiology*. 128 : 2157-2163.
- Comai L., Kosuge T., 1982. Cloning and Characterization of *iaaM*, a Virulence Determinant of *Pseudomonas savastanoi*. *Journal of Bacteriology*, 149: 40-46.
- Comai, L., and T. Kosuge. 1983. Transposable element that causes mutations in a plant pathogenic *Pseudomonas* sp. *J. Bacteriol.* 154:1162–1167.
- Cornara D., Cavalieri V., Dongiovanni C., Altamura G., Palmisano F., Bosco D., Porcelli F., Almeida R.P.P., Saponari M., 2016. Transmission of *Xylella fastidiosa* by naturally infected *Philaenus spumarius* (Hemiptera, Aphrophoridae) to different host plants. *J. Appl. Entomol.*, doi: 10.1111/jen.12365.
- Cornara D., Saponari M., Zeilinger A.R., de Stradis A., Boscia D., Loconsole G., Bosco D., Martelli G.P., Almeida R.P.P., Porcelli F., 2016. Spittlebugs as vectors of *Xylella fastidiosa* in olive orchards in Italy. *J. Pest. Sci.*, doi:10.1007/s10340-016-0793-0.
- Cornara, D., Loconsole, G., Boscia, D., Yokomi, R.K., Bosco, D., Porcelli, F., Saponari, M., Martelli, G. 2014. Survey of the auchenorrhyncha in the Salento Peninsula in search of putative vectors of *Xylella fastidiosa* Pauca Codiro strain. *Journal of Plant Pathology*. 96(4):S4.98.
- Corte A., Ciferri R., and Rui D., 1961. Infestazioni su ligustro da malformazioni fogliari dell'olivo. *Riv. Pat. Veg.*, Puvia S. III, 1 : 251-260.
- Cortese, C., Pérez-Martínez, I., Ramos, C., Buonauro, R., and Moretti, C. (2014). "The endophytes *Pantoea agglomerans* and *Erwinia oleae* colocalize with *Pseudomonas savastanoi* pv. *savastanoi* in the olive knots" in XX Convegno Nazionale Società di Patologia Vegetale, Environmentally loyal plant protection: from nano- to field-scale, 31.
- Costa, H. S., Raetz, E., Pinckard, T. R., Gispert, C., Hernandez-Martinez, R., Dumenyo, C. K., and Cooksey, D. A. 2004. Plant hosts of *Xylella fastidiosa* in and near southern California vineyards. *Plant Dis.* 88:1255-1261.
- Cronquist A., An Integrated System of Classification of Flowering Plants, Columbia University Press, 1981.
- Cruz, A. Braga da, & Tavares, R.M. (2005). Evaluation of programmed cell death in *Olea europaea* var. *Galega vulgar* suspension cell cultures elicited with *Pseudomonas savastanoi*, IX Congresso Luso-Espanhol de Fisiologia
- Vegetal, (September 2005), Available from <http://hdl.handle.net/1822/3437>
- Cutuli M., Campisi G., Marra F.P. and Caruso T. (2011). Vegetative growth and ecophysiological aspects in young olive plants inoculated with Olive leaf yellowing-associated virus (OLYAv). *Acta Italus Hortus*. 1, 356-361.
- da Clara Henriques M.I.E., (1994). Virus diseases of olive: an overlook. *Acta Hortic.* 356, 379-385 DOI: 10.17660/ActaHortic.1994.356.81
- Da Costa E Silva, O. C., Kosuge, T. 1991. Molecular characterization and expression analysis of the anthranilate synthase gene of *Pseudomonas syringae* subsp. *savastanoi*. *J. Bacteriol.* 173: 463-71.
- Davis M.J., Purcell A.H., Thompson S.V., 1978. Pierce's disease of grapevines: isolation of the causal bacterium. *Science* 199: 75-77.
- Djelouah K., Frasher D., Valentini F., D'onoghia A.M., Digiaro M., 2014. Direct tissue blot immunoassay for detection of *Xylella fastidiosa* in olive trees, *Phytopathologia Mediterranea* 53, 3, 559–564.
- Domenico, J. 1969. Relative infectivity of *Pseudomonas savastanoi* from olive and oleander. *Phytopathology*, 59: 1.
- dos Santos S.J., 2015. Transmissão de necrovírus de oliveira por *Olpidium virulentus*. Mestrado em Engenharia Agronómica. Universidade de évora escola de ciências e tecnologias departamento de fitotecnia. 71pp.
- Dye DW (1956) Oleander knot (*Pseudomonas savastanoi* (Erw. Smith) Stevens). *New Zealand Journal of Science and Technology* **38A**, 406–411.
- EFSA (European Food Safety Authority). 2015. Response to scientific and technical information provided by an NGO on *Xylella fastidiosa*. EFSA Journal, 13(4):4082 [13 pp.]
- El Air M., Mahfoudi N., Digiaro M., Najjar A., and Elbeaino T., (2011). Detection of Olive-infecting viruses in Tunisia. *Journal of Phytopathology*, 159(4), 201–238.
- El Beaino T., Saponari M., Minafra M.A., Castellano M.A., Savino V., Martelli G.P., 2005. Further characterization of Olive leaf yellowing-associated virus (*Olea europaea* L.). *Journal of plant pathology*. ISSN : 1125-4653.
- Elbeaino T., Digiaro M., De Stradis A., Martelli G.P., 2007. Identification of a second member of the family closteroviridae in mosaic-diseased figs. 89 (1): 119-124.
- Elbeaino T., Valentini F., Abou Kubaa R., Moubarak P., Yaseen T. and Digar M., 2014a. Multilocus sequence typing of *Xylella fastidiosa* isolated from olive affected by "olive quick decline syndrome" in Italy. *Phytopathologia Mediterranea* 53 (3): 533–542.
- Elbeaino T., Yaseen T., Valentini F., Ben Moussa I.E., Mazzoni V., D'ongh A.M., 2014b. Identification of three

- potential insect vectors of *Xylella fastidiosa* in southern Italy. *Phytopathologia Mediterranea*. 53 (1) : 328–332.
- Elbeaino T., Digiaro M., Martelli G.P., 2011. Complete sequence of Fig fleck-associated virus, a novel member of the family *Tymoviridae*. *Virus Research*. 161(2) : 198–202.
- Ercolani G.L., 1978. *Pseudomonas savastanoi* and other bacteria colonizing the surface of olive leaves in the field. *Journal of General Microbiology* 109: 245-257.
- Ercolani G.L., 1991. Distribution of epiphytic bacteria on olive leaves and the influence of leaf age and sampling time. *Microbial Ecology* 21: 35-48.
- Ercolani GL (1970) Presenza epifitica di *Pseudomonas savastanoi* (E. F. Smith) Stevens sull'olivo, in Puglia. *Phytopathologia Mediterranea* 10, 130–132.
- Ercolani GL (1985) Factor analysis of *Pseudomonas syringae* pv. *savastanoi* on the phylloplane of the olive. *Microbial Ecology* 11: 41–49.
- Ercolani, G. L. (1971). Presenza epifitica di *Pseudomonas savastanoi* (E. F. Smith) Stevens sull'Olivo, in Puglia. *Phytopathol. Mediterr.* 10, 130–132.
- Ercolani, G.L., 1983. Variability among isolates of *Pseudomonas syringae* pv. *savastanoi* from the phylloplane of the olive. *J. Gen. Microbiol.*, 129, 901–916.
- Erglmez S., Erkan S., 2014. Aydin, Balikesir ve Gzmir illerinde zeytin ağaçlarındaki viral hastalık etmenlerinin tanılanması ve bulunma durumlarının belirlenmesi. *Bitki Koruma Bülteni* 2014, 54(1):45-67.
- Essakhi S., Elbeaino T., Digiaro M., Saponari M., Martelli G.P., 2006. Nucleotide sequence variation in the HSP70h gene of Olive leaf yellowing associated virus. *Journal of Plant Pathology* 88: 285-291.
- Evidente A. and G. Surico, 1986. Isolation of indole-3-aldehyde from *Pseudomonas syringae* pv. *savastanoi*. *Journal of Natural Products* 49, 938–939 and references cited therein.
- Evidente A., G. Surico, N.S. Iacobellis and G. Randazzo, 1986. a-N-acetyl-indole-3-acetyl-e-L-lysine: a metabolite of indole-3-acetic acid from *Pseudomonas syringae* pv. *savastanoi*. *Phytochemistry* 25, 125–128.
- Grohmann F., "Oleaceae," Flora of Pakistan, vol. 59, p. 9, 1981.
- Fadel C., Digiaro M., Choueiri E., El Beaino T., Saponari M., Savino V., and Martelli G.P., 2005. On the presence and distribution of olive viruses in Lebanon. *Bulletin OEPP/EPPO Bulletin* 35, 33–36.
- Faggioli F., Barba M., 1995. A new elongated virus isolated from olive. *Acta Horticulturae* 386: 593-599.
- Faggioli F., Ferretti L., Pasquini G., Barba M., 2002. Detection of Strawberry latent ring spot virus in leaves of olive trees in Italy using one-step RT-PCR. *Journal of Phytopathology* 150: 636-639.
- Faggioli F., L. Ferretti, G. Albanese, R. Sciarroni, G. Pasquini, V. Lumia and M. Barba, 2005. Distribution of olive tree viruses in Italy as revealed by one-step RTPCR. *Journal of Plant Pathology* 87, 49–55.
- Feil, H., W. S. Feil, P. Chain, F. Larimer, G. DiBartolo, A. Copeland, A. Lykidis, S. Trong, M. Nolan, E. Goltsman, J. Thiel, S. Malfatti, J. E. Loper, A. Lapidus, J. C. Detter, M. Land, P. M. Richardson, N. C. Kyriides, N. Ivanova, and S. E. Lindow. 2005. Comparison of the complete genome sequences of *Pseudomonas syringae* pv. *syringae* B728a and pv. tomato DC3000. *Proc. Natl. Acad. Sci. USA* 102:11064–11069.
- Félix M.d.R., Varanda C.M.R., Clara M.I.E., 2011. Biology and molecular characterization of necroviruses affecting *Olea europaea* L.: a review. *Eur. J. Plant Pathol.*, 133:247–259
- Felix M.R and Clara M.I.E. (2006). Characterization of viruses occurring in *Olea europaea* L. In G.P. Rao, A, Valverde and C.I. Katis (Eds.). *Tech-niques in diagnosis of plant viruses* (pp. 173-216). Houston, TX, USA: Studium Press.
- Felix M.R., Cardoso J.M.S., Oliveira S. and Clara M.I.E. (2007). Biological and molecular charac-terization of *Olive latent virus 1*. *Plant Viruses* (pp. 170-177). Isleworth, UK: Global Science Books,
- Felix MRF, Clara MIE (2000) Two necrovirus with properties of *Olive latent virus 1* and of *Tobacco necrosis virus* from olive in Portugal. *Proc, 4<sup>th</sup> International Symposium on Olive Growing*, Bari, 2000. *Acta Horticult* 586: 725–728.
- Felix MRF, Louro TM, Clara MIE, Leit~ao F, Serrano J (2001) Preparation and application of an ELISA kit to identify *Olive latent virus 1* isolates obtained from *Olea europaea* L. cultivars grown in Portugal. *Proc, 11th Congress of the Mediterranean Phytopathological Union and 3rd Congress of the Sociedade Portuguesa de Fitopatologia, Evora*, 2001: 214–216.
- Félix, M. R., Cardoso, J. M. S., & Clara, M. I. E. (2006a). Single Strand Conformation Polymorphism analysis of Olive latent virus 1 isolates from *Olea europaea* L. *Proceedings of the Olivebioteq 2006- Second International Seminar "Biotechnology and Quality of Olive Tree Products around the Mediterranean Basin"*, Vol II, 243–246.
- Félix, M. R., Cardoso, J. M. S., Oliveira, S., & Clara, M. I. E. (2005a). Viral properties, primary structure and phylogenetic analysis of the coat protein of an Olive latent virus 1 isolate from *Olea europaea* L. *Virus Research*, 108, 195–198.
- Félix, M. R., Cardoso, J. M. S., Varanda, C. M. R., Oliveira, S., & Clara, M. I. E. (2005b). Complete nucleotide sequence of an Olive latent virus 1 isolate from olive trees. *Archives of Virology*, 150, 2403–2406.

- Ferretti L., Corsi B., Luongo L., Dal Cortivo C., Belisario A., 2017. A Survey of cherry leaf roll virus in intensively managed grafted english (persian) walnut trees in Italy. 99(2): 1-14.
- Félix, M. R., Clara, M. I., Leitão, F., & Serrano, J. (2002). Virus incidence in four *Olea europaea* cultivars evaluated by mechanical inoculation and immunological assays. Acta Horticulturae, 586(2), 721–724.
- Félix, M. R., Varanda, C., Cardoso, J. M. S., & Clara, M. I. E. (2006b). Plant root uptake of Olive latent virus 1 and Olive mild mosaic in single and mixed infections (pp. 516–517). Rhodes, Greece: Proceedings of XII MPU Congress.
- Félix, M.R., & Clara, M.I.E. (1998). Características biológicas e bioquímicas da estirpe G1, do Necrovirus Olive latent 1 isolado de *Olea europaea* L.. Actas da 2ª Reunião Bienal da Sociedade Portuguesa de Fitopatologia, pp 67.
- Felix, M.R., and Clara, M.I., 2002. Two necrovirus isolates with properties of olive latent virus 1 and of tobacco necrosis virus from olive in Portugal. Acta Hortic.586:725-728.
- Félix, M.R., Varanda, C.M.R., Cardoso, J.M.S., & Clara, M.I.E. (2004). Soil transmission of an olive isolate of Olive latent.
- Félix, M.R.F. (2007). Caracterização biomolecular de Olive latent virus 1 isolado de *Olea europaea* L. Tese de Doutoramento, Universidade de Évora, 114 p.
- Fernandes, A., and Marcelo, M. (2002). “A possible synergistic effect of *Erwinia* sp. on the development of olive knot symptoms caused by *Pseudomonas syringae* pv. *savastanoi* in *Olea europaea*” in Proceedings of the Fourth International Symposium on Olive Growing, Vols. 1 and 2, eds C. Vitagliano and G. P. Martelli (Valenzano: ISHS Acta Horticulturae), 729–731.
- Nguyen K.A., Förster H., Adaskaveg J.E., 2017. Quaternary Ammonium Compounds as New Sanitizers for Reducing the Spread of the Olive Knot Pathogen on Orchard Equipment. <https://doi.org/10.1094/PDIS-11-16-1578-RE>.
- Fichtner E., Kirkpatrick B., Ferguson L., Krueger W.H., DeBuse C., Zalom J., 2012. Epidemiology and management of olive knot disease: redefining management strategies based on pathogen population dynamics. California Olive Committee Final and Interim Research Reports. 1-4.
- Fichtner, E., Lightle, D., Krugner, R. 2015. Olive quick decline in Italy is associated with unique strain of *Xylella fastidiosa*. Subtropics newsletter. 13(1):2-3.
- Frisullo S., Camele I., Agosteo G.E., Boscia D., and Martelli G.P., 2014. Brief historical account of olive leaf scorch (“Brusca”) in the salento Peninsula of Italy and state-of-the-art of the olive quick decline syndrome. Journal of Plant Pathology, 96 (3) : 441-449.
- Gaffney, T., da Costa e Silva O., Yamada T., and Kosuge T., 1990. Indoleacetic acid operon of *Pseudomonas syringae* subsp. *savastanoi*: transcription analysis and promoter identification. J. Bacteriol. 172:5593-5601.
- Gallitelli D. and V. Savino, 1985. Olive latent virus 1. A single spherical virus isolated from olive in Apulia (southern Italy). Annals of Applied Biology 106, 295–303.
- García de los Ríos, J.E. (1989). Estudio acerca de la tuberculosis del olivo. Thesis (Ph. D.). Universidad Complutense de Madrid, Spain.
- García, A., Penyalver, R., & López, M.M. (2001). Sensibilidad de variedades de olivo a *Pseudomonas savastanoi* pv. *savastanoi*, causante de la tuberculosis. Fruticultura Profesional, No. 120, pp. 57-58, ISSN 1131-5660.
- Garcia, J.M. Yousfi, K. The postharvest of mill olives. Grasas Aceites. 57:16–24 (2006)
- Gardan L, Bollet C, Abu Ghorrah M, Grimont F, Grimont PAD (1992a) DNA relatedness among the pathovar strains of *Pseudomonas syringae* subsp. *savastanoi* Janse (1992a) and proposal of *Pseudomonas savastanoi* sp. nov. International Journal of Systematic Bacteriology 42, 606–612.
- Gardan L., David C., Morel M., Glickmann E., Abu-Ghorrah M., Petit A., Dessaix Y., Evidence for a correlation between auxin production and host plant species among strains of *Pseudomonas syringae* subsp. *savastanoi*, Appl. Environ. Microbiol. 58 (1992b) 1780–1783.
- Gaspar H.I.D., 2014. Estudo de um isolado de Olive latent virus 1: I. Caracterização molecular e II. Resposta à presença de bactérias promotoras de fitossanidade. Mestrado em Bioquímica, Dissertação. Universidade de Évora escola de ciências e tecnologia departamento de química. 145pp.
- Gaudin V., Vrain T., Jouanin L., 1994. Bacterial genes modifying hormonal balances in plants. Plant physiology and Biochemistry. 32(1): 11-29.
- Gemas V.J.V., Almadan M.C., Tenreiro R., Martins A., Fevereiro P., 2004. Genetic diversity in the Olive tree (*Olea europaea* L. subsp. *europaea*) cultivated in Portugal revealed by RAPD and ISSR markers. Genetic Resources and Crop Evolution 51: 501–511.
- Ghanney N., Locantore P., Nahdi S., Ferchichi A., Iacobellis N.S., 2016. Potential Biocontrol Effect of the Phylloplane Bacterium *Bacillus mojavensis* ABC-7 on the Olive Knot Disease. J. Plant Pathol Microbiol, 7:3.
- Giampetruzz A., Chiumenti M., Saponari M., Donvito G., Italiano A., Loconsole G., Boscia D., Cariddi C., Martelli G.P., Saldarellia P., 2015. Draft Genome

- Sequence of the *Xylella fastidiosa* CoDiRO Strain. *Genome Announcements*, 3 (1): 1-2.
- Giampetrucci A., Morelli M., Saponari M., Loconsole G., Chiumenti M., Boscia D., Savino V.N., Martelli G.P., Saldarelli P., 2016. Transcriptome profiling of two olive cultivars in response to infection by the CoDiRO strain of *Xylella fastidiosa* subsp. *Pauca*. *BMC Genomics* 2016;17:475 DOI: 10.1186/s12864-016-2833-9.
- Glass NL & Kosuge T (1986) Cloning of the gene for indoleacetic acid-lysine synthetase from *Pseudomonas syringae* subsp. *savastanoi*. *J Bacteriol* 166: 598–603.
- Glass NL & Kosuge T (1988) Role of indoleacetic acid-lysine synthetase in regulation of indoleacetic acid pool size and virulence of *Pseudomonas syringae* subsp. *savastanoi*. *J Bacteriol* 170: 2367–2373.
- Godena S., Bendini A., Giambanelli E., Cerretani L., Đermić D., Đermić E., 2012. Cherry leafroll virus: Impact on olive fruit and virgin olive oil quality. *European Journal of Lipid and Science Technology* 114: 535-541.
- Godena S., Damijanić K., Milotić A., 2009. Morphological characteristics of olive variety Rosinjola in Istria. *Pomologija Croatica*. 15(1-2): 27-36.
- Godena S., Dminić I., Đermić E., Silvana A.I., 2009. The occurrence of olive knot disease caused by *Pseudomonas savastanoi* in the northwest region of Istrian peninsula. *Zbornik predavanj in referatov 9. slovenskega posvetovanja o varstvu rastlin z mednarodno udeležbo* 467 Nova Gorica, 4 (5): 467-471.
- Godena S., Faggioli F., Luigi M., Saponari M., Loconsole G., Vonicina D. and Dermic' E., 2016. Incidence of viruses on autochthonous and introduced olive varieties in Croatian Istria detected by three diagnostic techniques. *Journal of Plant Pathology*. 98 (3): 657-660.
- Godini A., Mariani R., Pacifico A., Palasciano M., 2002. Repeatedly Reported but Hitherto Undescribed Olive Cultivars Native to Southern Italy. *Proc. 4<sup>th</sup> IS on Olive Growing* Eds. C. Vitagliano & G.P. Martelli *Acta Hort.* 586, ISHS 2002. 201-204.
- Gonzalez-Andujar, J.L., Expert system for pests, diseases and weeds identification in olive crops. *Expert Systems with Applications*. 36: 3278–3283. (2009).
- Gooch E., Ten Plus One Things You May Not Know about Olive, Epikouria Magazine, Triaina Publishers, 2005.
- Graniti A., 1990. Plant diseases in the mediterranean region. *Phytoparasitica* 18(1):57-65.
- Gratsia ME, Kyriakopoulou PE, Voloudakis AE, Fasseas C, Tzanetakis IE (2012) First Report of Olive mild mosaic virus and Sowbane mosaic virus in Spinach in Greece. *Plant Dis* 96: 1230–1230.
- Grieco F, Savino V, Martelli GP (1996a) Nucleotide sequence of the genome of a citrus isolate of olive latent virus 1. *Arch Virol* 141: 825–838.
- Grieco F., Alkowni R., Saponari M., Savino V., Martelli G.P., 2000. Molecular detection of olive viruses. *OEPP/EPPO, Bulletin Of PP/EPPO Bulletin* 30, 469-473
- Grieco, F., Parrella, G., & Vovlas, C. (2002). An isolate of olive latent virus 2 infecting castor bean in Greece. *Journal of Plant Pathology*, 84, 129–131.
- Grieco, F.; Dell'Orco, M.; Martelli, G. P.; 1996b: The nucleotide sequence of RNA1 and RNA2 of olive latent virus 2 and its relationships in the family Bromoviridae. *J Gen Virol* **77**, 2637–2644.
- Grieco, F.; Martelli, G. P.; Savino, V.; 1995: The nucleotide sequence of RNA3 and RNA4 of olive latent virus 2. *J Gen Virol* **76**, 929–937.
- Guan W., Shao J., Elbeaino T., Davis R.E., Zhao T., Huang Q., 2015. Specific Detection and Identification of American Mulberry-Infecting and Italian Olive-Associated Strains of *Xylella fastidiosa* by Polymerase Chain Reaction. *PLoS ONE* 10(6): e0129330. doi:10.1371/journal.pone.0129330
- Gutierrez F, Aranaud T, Albi AM. Influence of ecologic cultivation on virgin olive oil quality. *J Am Oil Chem Soc*. 76: 617–621 (1999).
- Haelterman R.M., Tolocka P.A., Roca M.E., Guzmán F.A., Fernández F.D., Otero M.L., 2015. First presumptive diagnosis of *Xylella fastidiosa* causing olive scorch in Argentina. *Journal of Plant Pathology*. 97 (2) : 393-393.
- Hagen K.S., 1963. Dependence of the Olive Fly, *Dacus oleae*, Larvae on Symbiosis with *Pseudomonas savastanoi* for the Utilization of Olive. *Nature* 209, 423 – 424.
- Haigen X., Sheng Q., Zhengmin H., Jianying G., Zongguo H., Hongying S., Shunping H., Hui D., Hairong W., Fanghao W., 2006. Biodiversity and Conservation (2006) 15:2893–2904.
- Hall BH, Cother EJ, Noble D, McMahon R, Wicks TJ (2003) First report of *Pseudomonas syringae* on olives (*Olea europaea*) in South Australia. *Australasian Plant Pathology* 32, 119–120. doi:10.1071/AP02053.
- Hall BH, Cother EJ, Whattam M, Noble D, Luck J, Cartwright D, 2004. First report of olive knot caused by *Pseudomonas savastanoi* pv. *savastanoi* on olives (*Olea europaea*) in Australia. *Australasian Plant Pathology* **33**, 433–6.
- Hartmann H.T., 2006. Olive Girdling Investigations seek to determine the best time and method for obtaining optimum fruit set. *California Agriculture*. 14-15.
- Hartmann, H.T. 1973. Chemicals to promote fruit abscission of olives. *Acta Hort.* 34:379–383.
- Hashemi-Tameh M., Bahar, Leila Zirak M., 'Candidatus Phytoplasma asteris' and 'Candidatus Phytoplasma aurantifolia', New Phytoplasma

- Species Infecting Apple Trees in Iran. *Journal of Phytopathology*. 162, Issue 7-8 August 2014 472–480
- Hashmi M.A., Khan A., Hanif M., Farooq U., Perveen S., 2015. Traditional Uses, Phytochemistry, and Pharmacology of *Olea europaea* (Olive). Hindawi Publishing Corporation Evidence-Based Complementary and Alternative Medicine. Article ID 541591, 29 pages.
- Hassani, D., Buonauro, R., and Tombesi, A. 2003. Response of some olive cultivars, hybrid and open pollinated seedling to *Pseudomonas savastanoi* pv. *savastanoi*. Pages 489-494 in: *Pseudomonas syringae* and Related Pathogens. Kluwer Academic Publishers, the Netherlands.
- Hatzopoulos P., Banilas G., Giannoulia K., Gazis F., Nikoloudakis N., Milioni D., Haralampidis K., 2002. Breeding, molecular markers and molecular biology of the olive tree. *Eur. J. Lipid Sci. Technol.* 104: 574–586.
- Henriques M.I.C., Leitao F., Potes M.F. and Serrano J.F., 1990. Virus detectados em *Olea europaea* L. Congresso Iberico de Ciencias horticolas, Lisboa pp. 245.
- Henriques M.I.C., Rei F.T., Leitão F.A., Serrano J.F. and Potes M.F. (1992). Virus diseases in *Olea europaea* L. cultivars. I. Immunodiagnosis of Strawberry latent ringspot virus. *Phytopathologia Mediterranea*. 31. 127-132.
- Hernandez-Martinez R., de la Cerda K.A., Costa H.S., Cooksey D.A., Wong F.P., 2007. Phylogenetic relationships of *Xylella fastidiosa* strains isolated from landscape ornamentals in Southern California. *Phytopathology* 97: 857-864.
- Hewitt WB, 1938. Leaf-scar infection in relation to the olive-knot disease. *Hilgardia* 12, 41–66.
- Hijazin R., Khlaif H., 2005. Detection of *Pseudomonas savastanoi* pv. *savastanoi* from olive and other hosts by polymerase chain reaction (PCR). *Dirasat. Agricultural Sciences* 32 (1), 71-77.
- Hopkins D.L., Mollenhauer H.H., 1973. Rickettia-like bacterium associated with Pierce's disease of grapes. *Science* 179: 298-300.
- Hopkins D.L., Purcell A.H., 2002. *Xylella fastidiosa*: cause of Pierce's disease of grapevine and other emergent diseases. *Plant Disease* 86: 1056-1066.
- Hopkins, D. L. (1989). *Xylella fastidiosa*: xylem-limited bacterial pathogen of plants. *Annual Review of Phytopathology*, 29, 271–290.
- Hopkins, D. L. (2014). Control strategies for *Xylella fastidiosa*. International Symposium on the European Outbreak of *Xylella fastidiosa* in Olive, *Journal of Plant Pathology*, 96, S4.99.
- Horne W.M., Parker W.B., and Daines L.L., 1912: The method of spreading of the olive knot disease. *Phytopathology* 2, 101—105.
- Montano, H.G., Davis, R.E., Dally, E.L., Hogenhout, S., Pimental, J.P., and Brioso, P. ST.2001. 'Candidatus phytoplasma brasiliense', a new phytoplasma taxon associated with hibiscus witches' broom disease. *Int. J. Syst. and Evol. Microbiol.* 51:1109-1118
- Hosni, T. (2010). Interaction Between *Pseudomonas savastanoi* pv. *savastanoi*, the Causal Agent of Olive Knot, and the Endophytic Bacterial Species Associated With the Knot. Ph.D. thesis. University of Perugia, Perugia, Italy.
- Hosni, T., Moretti, C., Devescovi, G., Suarez-Moreno, Z. R., Barek Fatmi, M., Guarnaccia, C., et al. (2011). Sharing of quorum-sensing signals and role of interspecies communities in a bacterial plant disease. *ISME J.* 5, 1857–1870. doi: 10.1038/ismej.2011.65.
- Hutcheson, S.W., and Kosuge T., 1985. Regulation of 3-indoleacetic acid production in *Pseudomonas syringae* pv. *savastanoi*. *J. Biol. Chem.* 260:6281-6287.
- Hutzinger, O., and T. Kosuge. 1968. 3-Indoleacetic-L-lysine, a new conjugate of 3-indoleacetic acid produced by *Pseudomonas savastanoi*, p. 183–194. In F. Wightman and G. Setterfield (ed.), *Biochemistry and physiology of plant growth substances*. The Runge Press, Ltd., Ottawa, Ontario, Canada.
- Hutzinger, O., and T. Kosuge. 1968. 3-Indoleacetic-L-lysine, a new conjugate of 3-indoleacetic acid produced by *Pseudomonas savastanoi*, p. 183–194. In F. Wightman and G. Setterfield (ed.), *Biochemistry and physiology of plant growth substances*. The Runge Press, Ltd., Ottawa, Ontario, Canada.
- Iacobellis N.S., Caponero A. and Evidente A., 1998. Characterization of *Pseudomonas syringae* ssp. *savastanoi* strains isolated from ash. *Plant Pathology* (1998) 47, 73–83.
- Iacobellis NS, Sisto A and Surico G (1993) Occurrence of unusual strains of *Pseudomonas syringae* subsp. *savastanoi* on olive in central Italy. EPPO Bulletin 23: 429–435.
- EL-Morsy Sh.I., EL-Sheikh M.A., Abd El-Razik R.A., Youssef S.A., Shalaby A.A., 2017. Molecular Identification of Strawberry Latent Ring Spot Virus (SLRSV) In Egypt. *Journal of Basic and Environmental Sciences* 4 (2017) 24-33.
- Iacobellis NS, Sisto A, Surico G, Evidente A, Di Maio E. 1994. Pathogenicity of *Pseudomona syringae* subsp. *savastanoi* mutants defective in phytohormone production. *Journal of Phytopathology* 140: 238–248.
- Iacobellis NS. 2001. Olive knot. In: Maloy OC, Murray TD, eds. *Encyclopedia of plant pathology*. New York, NY, USA: John Wiley & Sons, 714–715.
- Iacobellis, N. S., Contesini A. M., and Surico G., 1995. Bacteriocin production by *Pseudomonas syringae* subsp. *savastanoi*. *Phytopathol. Mediterr.* 34: 15–22.

- Janse J.D. and Obradovic A., 2010. *Xylella fastidiosa*: its biology, diagnosis, control and risks. *Journal of Plant Pathology*, 92 (1): 35- 48.
- Janse J.D., 1991: Pathovars of *Pseudomonas syringae* subsp. *savastanoi*: discrimination on the basis of whole cell fatty acids. In: Proc. 4th Working Group on *Pseudomonas syringae* pathovoras, Firenze, pp. 337-339.
- Janse, J.D. (1982). *Pseudomonas syringae* subsp. *savastanoi* ex Smith subsp. nov., nom. rev., the bacterium causing excrescences on Oleaceae and Nerium oleander L. *International Journal of Systematic Bacteriology*, Vol. 32, No. 2, (April 1982), pp. 166-169, ISSN: 1466-5026.
- Jardak T., Jerraya A. et Mahjoub M., 2004. La protection intégrée de l'oléiculture dans les pays de l'Afrique du Nord. FAO, Bureau sous régional pour l'Afrique du Nord, SNEA-Tunis. Tunisie, 69-71.
- Jiménez Diaz, R.M. (1985) Les maladies de l'olivier. *Olivae* 2(8), 24–28.
- Kacem M., Kazouz F., Merabet C., Rezki M., De Lajudie P., Bekki A., 2009. Antimicrobial activity of *Rhizobium* sp. strains against *Pseudomonas savastanoi*, the agent responsible for the olive knot disease in Algeria. *Grasas y Aceites* 60: 139-146.
- Kailis G., Sweeney S., 2002. Olives in Australia. Proc. 4th IS on Olive Growing Eds. C. Vitagliano & G.P. Martelli Acta Hort. 385-388.
- Kanakis P., A. Termentzi, T. Michel, E. Gikas, M. Halabalaki, and A.-L. Skaltsounis, "From olive drupes to olive Oil. An HPLC-orbitrap-based qualitative and quantitative exploration of olive key metabolites," *Planta Medica*, vol. 79, no. 16, pp. 1576– 1587, 2013.
- Kanematsu S, TagaY, Morikawa T (2001) Isolation of Olive latent virus 1 from tulip in Toyama Prefecture. *J Gen Plant Pathol* 67: 333–334
- Kaniewski D., E. van Campo, T. Boiy, J.-F. Terral, B. Khadari, and G. Besnard, "Primary domestication and early uses of the emblematic olive tree: palaeobotanical, historical and molecular evidence from the Middle East," *Biological Reviews*, vol. 87, no. 4, pp. 885–899, 2012.
- Kavak H. and Üstün N., 2009. Oleander knot caused by *Pseudomonas savastanoi* pv. *nerii* in Turkey. *Journal of Plant Pathology*, Vol. 91, No. 3 (November 2009), pp. 701-703.
- Kerenyi A., Bihary D., Venturi V., Pongor S., 2013. Stability of Multispecies Bacterial Communities: Signaling Networks May Stabilize Microbiomes. *Microbiome Stability*, 8 (3): 1-10.
- Khlaif H., 2006. Olive Knot Disease in Jordan. National Center for Agricultural Research & Extension. Homepage: <http://www.ncartt.gov.jo/>. 2 (4).
- Kiritsakis, A. Olive oil: from the tree to the table, 2nd ed. Trumbull (CT): Food and Nutrition Press (1998).
- Kosuge, T.M., Heskett G., and Wilson E.E., 1966. Microbial synthesis and degradation of indol-3-acetic acid. I. The conversion of L-tryptophan to indol-3-acetamide by an enzyme system from *Pseudomonas savastanoi*. *J. Biol. Chem.* 241:3738-3744.
- Krid S, Rhouma A, Queseda JM, Penyalver R, Gargouri A (2009) Delineation of *P. savastanoi* pv. *savastanoi* strains isolated in Tunisia by ramdom-amplified polymorphic DNA analysis. *J Appl Microbiol* 106:886–894
- Krid S., Bouaziz M., Triki M.A., Gargouri A. and Rhouma A., 2011b. Inhibition of olive knot disease BY polyphenols extracted from olive mill waste water. *Journal of Plant Pathology*. 93(3): 561-568.
- Krid S., Gharsallaoui M., Triki M.A., Rhouma A., 2011a. Epidemiologie et etiologie de *Pseudomonas savastanoi* pv. *savastanoi*, agent causal de la tuberculose de l'olivier en Tunisie. *Revue Ezzaitouna* 12 (1) : 1-10.
- Krid S., Rhouma A., Mogou I., Quesada J.M., Nesme X., and Gargouri A., 2010. *Pseudomonas savastanoi* endophytic bacteria in olive tree knots and antagonistic potential of strains of *Pseudomonas fluorescens* and *Bacillus subtilis*. *Journal of Plant Pathology*, 92 (2), 335-341.
- Krid S., Triki M.A., Gargouri A., Rhouma A., 2012. Biocontrol of olive knot disease by *Bacillus subtilis* isolated from olive leaves. *Ann Microbiol.* 62:149–154.
- Krueger W.H., Tevitdale B.L., Scroth M.N., Metzidakis I.T., Voyatzis D.G., 1999. Improvements in the control of olive knot disease. *Acta Horticulturae* 474 (2): 567-571.
- Krueger WH, Teviotdale BL, Schroth MN, Metzidakis IT, Voyatzis DG (1999) Improvements in the control of olive knot disease. Proceedings of the 3<sup>rd</sup> International Symposium on Olive Growing. Chania, Crete, Greece, 22–26 September 1997, Vol. 2. (Eds IT Metzidakis, DG Voyatzis). *Acta Horticulturae* **474**, 567–571.
- Krugner R., Johnson M.W., Chen J., 2010. Evaluation of pathogenicity and insect transmission of *Xylella fastidiosa* strains to olive plants. California Olive Committee. Final Report 2010: <http://calolive.org/wp-content/uploads/Research-Reports-2010.pdf>.
- Krugner, R. 2010. Evaluation of pathogenicity and insect transmission of *Xylella fastidiosa* strains to olive plants [abstract]. California Olive Committee Annual Research Report. P. 48-56.
- Krugner, R., Sisterson, M. S., Chen, J., Stenger, D. C., and Johnson, M. W. 2014. Evaluation of olive as a host of *Xylella fastidiosa* and associated sharpshooter vectors. *Plant Dis.* 98:1186-1193.
- Kuo, T.T., and T. Kosuge. 1969. Factors influencing the production and further metabolism of indole-3-acetic acid

- by *Pseudomonas savastanoi*. J. Gen. Appl. Microbiol. 15:51-63.
- Lamichhane J.R., Varvaro L., (2013). Epiphytic *Pseudomonas savastanoi* pv. *savastanoi* can infect and cause olive knot disease on *Olea europaea* subsp. *Cuspidate*. Australasian Plant Pathol. 42:219–225.
- Lang, G.A. 1987. Physiological studies of ethylene-induced olive (*Olea europaea* L.) fruit and leaf abscission. PhD diss. Univ. of California, Davis.
- Lavermicocca P., Lonigro S.L., Valerio F., Evidente A., Visconti A., 2002. Reduction of olive knot disease by a bacteriocin from *Pseudomonas syringae* pv. *ciccaronei*. Applied and Environmental Microbiology 68: 1403-1407.
- Lavermicocca P., Valerio F., Lonigro S.L., Lazzaroni S., Evidente A., Visconti A., 2002. Control of Olive Knot Disease with a Bacteriocin. Kluer Academic publishers. 451-457.
- Lavermicocca, P., M. Gobbetti, A. Corsetti, and L. Caputo. 1998. Characterization of lactic acid bacteria isolated from olive phylloplane and table olive brines. Ital. J. Food Sci. 10:27–39.
- Lavermicocca, P., M. Gobbetti, A. Corsetti, and L. Caputo. 1998. Characterization of lactic acid bacteria isolated from olive phylloplane and table olive brines. Ital. J. Food Sci. 10:27–39.
- Lavermicocca, P., S. L. Lonigro, A. Evidente, and A. Andolfi. 1999. Bacteriocin production by *Pseudomonas syringae* pv. *ciccaronei* NCPPB2355. Isolation and partial characterization of the antimicrobial compound. J. Appl. Microbiol. 86:257–265.
- Leitao, F., Serrano, J.F., Fatima Potes, M., Ivone Clara, M., Rosario Felix, M., Rei, F. and Lopes, J. (1999). Preliminary results on clonal and sanitary selection of *Olea europaea* L.CV. 'SANTULHANA' in north-east of Portugal. Acta Hortic. 474, 163-166.
- Lipshitz N., R. Gophna, M. Hartman, and G. Biger, "The beginning of olive (*Olea europaea*) cultivation in the old world: a reassessment," Journal of Archaeological Science, vol. 18, no. 4, pp. 441–453, 1991.
- Lobão, D.L., Félix, M.R., Clara, M.I.E., Oliveira, S., Leitão, F. A., & Serrano, J.F. (2002). Detection of Olive latent virus 1 in *Olea europaea* L. tissues by reverse transcriptionpolymerase chain reaction. Book of Abstracts XIII Congresso Nacional de Bioquímica, Lisboa, 102 p.
- Loconsole G., Potere O., Boscia D., Altamura G., Djelouah K., Elbeaino T., Frasher D., Lorusso D., Palmisano F., Pollastro P., Silletti M.R., Trisciuzzi N., Valentini F., Savino V. and Saponari M., 2014a. Detection of *Xylella fastidiosa* in olive trees by molecular and serological methods. Journal of Plant Pathology, 96 (1), 7-14.
- Loconsole G., Potere O., Elbeaino T., Frasher D., Frisullo S., Palmisano F., Boscia D., Saponari M., 2014b. Interlaboratory validation of molecular and serological diagnosis of *Xylella fastidiosa* strain CoDiRO in susceptible host plants.
- Loconsole G., Saponari M., Boscia D., D'Attoma G., Morelli M., Martelli G.P., Almeida R.P.P., 2016. Intercepted isolates of *Xylella fastidiosa* in Europe reveal novel genetic diversity. Eur. J. Plant Pathol., DOI 10.1007/s10658-016-0894-x
- Loconsole G., Saponari M., Bottalico G., Faggioli F., Albanese G. and Savino V., 2013. Production of Nuclear Stock olive plants for certification programs. La rivista DI scienza dell'alimentazione, numero 1, Gennaio-Marzo 2013, Anno 42. 75-84.
- Loconsole, G., Saponari, M., Faggioli, F., Albanese, G., Bouyahia, H., Elbeaino, T., Materazzi, A., Nuzzaci, M., Prota, V., Romanazzi, G., Trisciuzzi, N., & Savino, V. (2010). Inter-laboratory validation of PCR-based protocol for detection of olive viruses. Bulletin OEPP/EPPO, 40, 423–428.
- Lopes J.R.S., Landa B.B., and Fereres A., 2014. A survey of potential insect vectors of the plant pathogenic bacterium *Xylella fastidiosa* in three regions of Spain. Spanish Journal of Agricultural Research 2014 12(3): 795-800.
- Lopez-Villalta MC (1999) *Pseudomonas savastanoi* SMITH. In 'Olive pest and disease management'. pp. 175–178. (IOOC: Madrid, Spain).
- Luigi M., Godena S., Đermić E., Barba M. and Faggi F., 2011. Detection of viruses in olive trees in Croatian Istria. Phytopathol. Mediterr., 50 : 150–153.
- Luigi M., Manglli A., Thomaj F., Buonauro R., Barba M., Faggioli F., 2009. Phytosanitary evaluation of olive germplasm in Albania. Phytopathol. Mediterr., 48: 280–284
- Macdonald, E. M. S., G. K. Powell, D. A. Regier, N. L. Glass, F. Roberto, T. Kosuge, and R. O. Morris. 1986. Secretion of zeatin, ribosylzeatin, and ribosyl-1-methylzeatin by *Pseudomonas savastanoi*: plasmid-coded cytokinin biosynthesis. Plant Physiol. 82:742–747.
- Magie A.R., Wilson E.E., Expressions of virulence among isolates of *Pseudomonas savastanoi* from olive and oleander, Proc. Am. Phytopathol. Soc., Phytopathology 52 (1962) 741 (Abst.).
- Lee, I.M, Dawn, E, Gundersen-Rindal, D.E, Davis, R.E., and Bartoszyk, M., 1998. Revised classification scheme of phytoplasmas based on RFLP analyses of 16S rRNA and ribosomal protein gene sequences. Int. J. Syst. Bacteriol. 48:1153-1169.
- Mahhou A., Nabil Y., Hadiddou A., Oukabli A. and Mamouni A., 2012. Performance of the Arbequina, Haouzia and Menara olive varieties in rainfed conditions in the Meknès region of Morocco. Official magazine of the International Olive Counc.3-21.

- Maldonado-González, M., Prieto, P., Ramos, C., and Mercado-Blanco, J. (2013). From the root to the stem: interaction between the biocontrol root endophyte *Pseudomonas fluorescens* PICF7 and the pathogen *Pseudomonas savastanoi* NCPPB 3335 in olive knots. *Microb. Biotechnol.* 6, 275–287.
- Mang S.M., Frisullo S., Elshafie H.S. and Camele I., 2016. Diversity Evaluation of *Xylella fastidiosa* from Infected Olive Trees in Apulia (Southern Italy). *Plant Pathol.* J. 32(2) : 102-111.
- Marcelo, A., Fernandes, M., Fatima Potes, M., and Serrano, J. F. 1999. Reactions of some cultivars of *Olea europaea* L. to experimental inoculation with *Pseudomonas syringae* pv. *savastanoi*. *Acta Hortic.* 474:581-584.
- Marchi G., Mori B., Pollacci P., Mencuccini M., Surico G., 2009. Systemic spread of *Pseudomonas savastanoi* pv. *savastanoi* in olive explants *Plant Pathology* 58: 152-158.
- Marchi G., Sisto A., Cimmino A., Andolfi A., Cipriani M.G., Evidente A., Surico G., 2006. Interaction between *Pseudomonas savastanoi* pv. *savastanoi* and *Pantoea agglomerans* in olive knots. *Plant Pathology* 55: 614-624.
- Marchi G., Viti C., Giovannetti L., Surico G., 2005. Spread of levan-positive populations of *Pseudomonas savastanoi* pv. *savastanoi*, the causal agent of olive knot, in central Italy *European Journal of Plant Pathology* 112: 101-112.
- Maretti G.P., Gallitelli D., 1985. virosi dell'olivo. *Italia agricola*. 122 (2) : 150-156.
- Marlow J.L., Kosuge T., 1972. Tryptophan and Indoleacetic Acid Transport in the Olive and Oleander Knot Organism *Pseudomonas savastanoi* (E. F. Smith) Stevens.
- Martelli G.P., 1998. Infectious diseases and olive tree certification: general panorama. <http://agris.fao.org/agris-search/search.do?recordID=ES1998001911>
- Martelli G.P., 2013a. Disseccamento rapido dell'olivo. *Georgofili INFO*, 30 Ottobre 2013. ([www.georgofili.info](http://www.georgofili.info)).
- Martelli G.P., 2013b. A Brief Outline of Infectious Diseases of Olive. *Palestine Technical University Research Journal*, 2013, 1(1), 01-09.
- Martelli G.P., 2016. The current status of the quick decline syndrome of olive in southern Italy. *Phytoparasitica*. 44:1–10.
- Martelli G.P., Boscia D., Porcelli F., Saponari M., 2016. The olive quick decline syndrome in south-east Italy: a threatening phytosanitary emergency. *Eur. J. Plant Pathol.*, 144:235–243.
- Martelli G.P., Salerno M., Savino V. Prota U., 2002. An Appraisal of Diseases and Pathogens of Olive. *Proc. 4th IS on Olive Growing* Eds. C. Vitagliano & G.P. Martelli. *Acta Hort.* 586, ISHS 2002. 701-708.
- Martelli G.P., Savino V., Di Terlizzi B., Catalano L., Sabanadzovic S. and Greco N. (1995). Viruses and certification of olive in Apulia (Southern Italy). *Acta Horticulturae*. 386, 569-573.
- Martelli GP (2002) Certification of olive: the Italian experience. Séminaire international sur l'oléiculture: acquis de recherche et contraintes du secteur. Marrakech (MA).
- Martelli, G. P., Sabanadzovic, S., Savino, V., Abu-Zurayk, A. R., & Masannat,M. (1995). Virus-like disease and viruses of olive in Jordan. *Phytopathologia Mediterranea*, 34, 133–136.
- Martelli, G. P., Yilmaz, M. A., Savino, V., Baloglu, S., Grieco, F., Guldur, M. E., Greco, N., & Laforteza, R. (1996). Properties of a citrus isolate of olive latent virus 1, a new necrovirus. *European Journal of Plant Pathology*, 102, 527–536.
- Martelli, G.P. 1999. Infectious diseases and certification of olive: an overview. *EPPO Bull.* 29:127-133.
- Martelli, G.P., Agranovsky, A.A., Bar-Joseph, M., Boscia, D., Candresse, T., Coutts, R.H.A., Dolja, V.V., Falk, B.W., Gonsalves, D., Jelkmann, W., Karasev, A.V., Minafra, A., Namba, S., Vetten, H. J., Wisler, G.C., and Yoshikawa, N., 2002. The family Closteroviridae revised. *Arch. Virol.* 147:2039-2044.
- Martin, G.C. 1986. Olive harvest in California, United States of America. *Olivae* 3:11-20.
- Martos-Moreno C., Lopez-Escudero F.J., Blanco-Lopez M.A., 2006. Resistance of Olive cultivars to the defoliating pathotype of *Verticillium dahliae*. *HortScience*. 41 (5): 1313-1316.
- Matas I.M., Pérez-Martínez I., Quesada J.M., Rodríguez-Herva J.J., Penyalver R., Ramos C., 2009. *Pseudomonas savastanoi* pv. *savastanoi* contains two *iaaL* paralogs, one of which exhibits a variable number of a trinucleotide (TAC) tandem repeat. *Applied and Environmental Microbiology* 75: 1030-1035. 15-822.
- Matas IM, Castaneda-Ojeda MP, Aragon IM, Antunez-Lamas M, Murillo J, Rodriguez-Palenzuela P, Lopez-Solanilla E & Ramos C (2014) Translocation and functional analysis of *Pseudomonas savastanoi* pv. *savastanoi* NCPPB 3335 type III secretion system effectors reveals two novel effector families of the *Pseudomonas syringae* complex. *Mol Plant Microbe Interact* doi: 101094/MPMI-07-13-0206-R.
- Matas IM, Lambertsen L, Rodriguez-Moreno L & Ramos C (2012) Identification of novel virulence genes and metabolic pathways required for full fitness of *Pseudomonas savastanoi* pv. *savastanoi* in olive (*Olea europaea*) knots. *New Phytol* 196: 1182–1196.

- Matas, I.M. (2010). Genómica funcional de la interacción *Pseudomonas savastanoi* pv. *savastanoi*- olivo. Thesis (Ph. D.). Universidad de Málaga, Spain.
- Materazzi A., Toni S., Panettoni A., Osti M., Triolo E., 1996. On the presence of a new isometric virus in *Olea europaea* L.. In: *Proceedings Convegno Annuale della Società Italiana di Patologia Vegetale, Udine* 1996, 57-59.
- Mehri, H., Mehri-Kamoun, R., Hibar, K., 2009. Response of 'Ousleti' Olive Pollen to *Pseudomonas syringae* pv. *Savastanoi* Culture Filtrate. The African Journal of Plant Science and Biotechnology. 3(1): 44-48.
- Merciega V, Boscia D, Savino V (1996) Comparison of five isolates of olive latent virus1. Phytopathol Medit 35: 1–8.
- Mirik M., Aysan Y., 2011. Phenotypic and genotypic characterization of *Pseudomonas savastanoi* pv. *savastanoi* isolates and disease prevalence of Olive Knot Disease in Marmara Region of Turkey. Journal of Agricultural Sciences (Turkey), 2011 - agris.fao.or
- Mirik M., Aysan Y., Sahin F., 2006. Isolation, pathogenicity and characterization of *Pseudomonas savastanoi* pv. *savastanoi* from different hosts in Turkey. Proceedings of the 12<sup>th</sup> Congress of the Mediterranean Phytopathological Union, Rhodes2 006:3 21-325.
- Moffett ML (1983) Bacterial plant pathogens recorded in Australia. In 'Plant bacterial diseases'. (Eds PC Fahy, GJ Persley) pp. 317–336. (Academic Press, Australia).
- Montemurro, C., Simeone, R., Blanco, A., Saponari, M., Bottalico, G., Savino, V., Martelli, G.P. and Pasqualone, A. (2008). Sanitary selection and molecular characterization of olive cultivars grown in Apulia. Acta Hortic. 791, 603-609
- Moral, J. De La Rosa, R. Leon, L. Barranco, D. Michailides, T.J. Trapero, A. High susceptibility of olive cultivar FS-17 to *Alternaria alternata* in Southern Spain. Plant Dis 92:1252 (2008a).
- Morea A., M., Zaccaro, F., Palumbo, G., and Iacobellis, N.S. (1999) Isolation and characterization of *Pseudomonas syringae* susp. *savastanoi* mutants defective in hypersensitive response elicitation and pathogenicity. *J Phytopathol* **147**: 321–330.
- Moretti C., Hosni T., Vandemeulebroecke K., Brady C., De Vos P., Buonauro R. and Cleenwerck I., 2011. *Erwinia oleae* sp. nov., isolated from olive knots caused by *Pseudomonas savastanoi* pv. *Savastanoi*. International Journal of Systematic and Evolutionary Microbiology, 61, 2745–2752.
- Moretti, C., Cortese, C., Passos da Silva, D., Devescovi, G., Torelli, E., Venturi, V., et al. (2014a). "Draft genome sequence of a hypersensitive reaction-inducing *Pantoea agglomerans* strain isolated from olive knots caused by *Pseudomonas savastanoi* pv. *Savastanoi*," in 13th International Conference on Plant Pathogenic Bacteria (Shanghai, China), S1–P2.
- Moretti, C., Cortese, C., Passos da Silva, D., Venturi, V., Firrao, G., and Buonauro, R. (2014b). Draft genome sequence of *Erwinia oleae*, a bacterium associated with olive knots caused by *Pseudomonas savastanoi* pv. *savastanoi*. Genome Announc. 2, e01308-14.
- Moretti, C., Cortese, C., Passos da Silva, D., Venturi, V., Ramos, C., Firrao, G., et al. (2014c). Draft genome sequence of *Pseudomonas savastanoi* pv. *savastanoi* strain DAPP-PG722, isolated in Italy from an olive plant affected by knot disease. Genome Announc. 2, e00864 14.
- Moretti, C., Cortese, C., Passos da Silva, D., Venturi, V., Torelli, E., Firrao, G., et al. (2014d). Draft genome sequence of a hypersensitive reaction-inducing *Pantoea agglomerans* strain isolated from olive knots caused by *Pseudomonas savastanoi* pv. *savastanoi*. Genome Announc. 2, e00774-14.
- Moretti, C., Ferrante, P., Hosni, T., Valentini, F., D'Onghia, A., Fatmi, M.B., Fatmi, M.B. and Buonauro, R. (2008) Characterization of *Pseudomonas savastanoi* pv. *savastanoi* strains collected from olive trees in different countries. In: *Pseudomonas syringae* Pathovars and Related Pathogens—Identification, Epidemiology and Genomics (Fatmi, M.B., Collmer, A., Iacobellis, N., Mansfield, J., Murillo, J., Schaad, N., et al., eds), pp. 321–329. Dordrecht: Springer Netherlands.
- Moukhli A., 1984. Olive reactions to the olive knot and identification of the causal agent (*Pseudomonas syringae* subsp. *savastanoi* pv. *Olea*). Direction de la programmation et des affaires économiques, Ministère de l'agriculture, du développement rural et des pêches maritimes. <http://www.agriculture.gov.ma/>
- Mourad K., Kazouz H., 2009. Properties of bacteriocin like substances produced by *Rhizobium* sp. ORN83 and ORN24 strains previously isolated in saline soils of Algeria. International Journal of Mediterranean Ecology. 23: 91-98.
- Moussa I.E.B., Mazzoni V., Valentini F., Yaseen T., Lorusso D., Speranza S., Digiaro M., Varvaro L., Krugner R., D'Onghia A.M., 2016. Seasonal Fluctuations of Sap-Feeding Insect Species Infected by *Xylella fastidiosa* in Apulian Olive Groves of Southern Italy. Journal of Economic Entomology, 109(4) : 1512–1518.
- Mugnai L., Giovannetti L., Ventura S., Surico G., 1994. The Grouping of Strains of *Pseudomonas syringae* subsp. *Savastanoi* by DNA Restriction Fingerprinting. J. Phytopathology 142,209—218.
- Noble RJ, Hynes HJ, McCleery FC, Brimingham WA (1935). Plant diseases recorded in New South Wales. New South Wales Department of Agriculture Science Bulletin No. 46.

- Oliveira, M. R., & Luz, J. P. (2003). Susceptibilidade varietal de cultivares de oliveira a *Pseudomonas savastanoi* na região da Beira Interior. (Poster presented at the III Simpósio Nacional de Olivicultura, Escola Superior Agrária de Castelo Branco, Castelo Branco, Portugal).
- Orešek E., Seljak G., 2015. *Xylella fastidiosa* - new threat for Slovenian olive growing. Zbornik predavanj in referatov 12. Slovenskega posvetovanja o varstvu rastlin z mednarodno udeležbo, 3.-4. marec 2015, Ptuj, Slovenija. pp.28-32 ref.8
- Osman WA, Tarabeih AM, Michail SH (1980a) Studies on the distribution of olive knot disease induced by *Pseudomonas savastanoi* in Iraq. *Mesopotamia Journal of Agriculture* **15**, 235–243.
- Osman WA, Tarabeih AM, Michail SH (1980b). Studies on olive knot disease in Iraq with reference to response of different cultivars. *Mesopotamia Journal of Agriculture* **15**, 245–261
- Ouzari H., Khsairi A., Raddadi N., Jaoua L., Hassen A., Zarrouk M., Daffonchio D., Boudabous A., 2008. Diversity of auxin-producing bacteria associated to *Pseudomonas savastanoi*-induced olive knots. *Journal of Basic Microbiology* 48: 1-8.
- Pacini E & Cresti M (1977) Viral particles in developing pollen grains of *Olea europaea*. Planta 137, 1-4.
- Palm, C.J., Gaffney, T., Kosuge, T. 1989. Cotranscription of gene encoding indoleacetic acid production in *Pseudomonas savastanoi*. 1. Bacteriol. 171: 1002-9.
- Palumbo, J.D. Baker, J.L. Mahoney, N.E. Isolation of bacterial antagonists of *Aspergillus flavus* almonds. *Microbial Ecol.* 52(1): 45- 52 (2006)
- Panagopoulos CG (1993) Olive knot disease in Greece. *Bulletin OEPP. EPPO Bulletin. European and Mediterranean Plant Protection Organisation* **23**, 417–422.
- Pantaleo, V., Grieco, F., Castellano, M. A., & Martelli, G. P. (1999). Synthesis of infectious transcripts of olive latent virus 1: genes required for RNA replication and virus movement. *Archives of Virology*, 144, 1071–1079.
- Pantaleo, V., Grieco, F., Di Franco, A., & Martelli, G.P. (2006). The role of the C-terminal region of Olive latent virus 1 coat protein in the host systemic infection. *Archives of Virology* 151, 1973–1983
- Pantaleo, V., Saponari, M., & Gallitelli, D. (2001). Development of a nested PCR protocol for detection of olive infecting virus in crude extracts. *Journal of Plant Pathology*, 83(2), 143–146.
- Pashaei R., 2016. Description of Chaperone Protein or Q9PEQ0 (HTPG\_XYLFA) in *Xylella Fastidiosa* Organism byBLAST, Jpred4, and FTMap methods with help of Pymol Software. *Int J Genomics Proteomics Metabolomics Bioinformatics*. 1(3), 14-16.
- Passos da Silva D., G. Devescovì, Paszkiewicz K., Moretti C., Buonauro R., Studholme D.J., Venturia V., 2013. Draft Genome Sequence of *Erwinia toletana*, a Bacterium Associated with Olive Knots Caused by *Pseudomonas savastanoi* pv. *Savastanoi*. 1(3):e00205-13. doi:10.1128/genomeA.00205-13.
- Passos da Silva, D., Pilar Castaneda-Ojeda, M., Moretti, C., Buonauro, R., Ramos, C., and Venturi, V. (2014). Bacterial multispecies studies and microbiome analysis of a plant disease. *Microbiology* 160, 556–566.
- Penyalver, R., A. García, A. Ferrer, E. Bertolini, and M. M. Lopez. 2000. Detection of *Pseudomonas savastanoi* pv. *savastanoi* in olive plants by enrichment and PCR. *Appl. Environ. Microbiol.* 66:2673–2677.
- Penyalver, R., García, A., Del Río, C., Caballero, J.M., Pinochet, J., Piquer, J., & López, M.M. (2003). Sensibilidad varietal del olivo a la tuberculosis causada por *Pseudomonas savastanoi* pv. *savastanoi*. Agrícola vergel: Fruticultura, horticultura, floricultura, No. 253, pp. 13-17, ISSN 0211-2728.
- Penyalver, R., García, A., Ferrer, A., & López, M.M. (1998). La tuberculosis del olivo: diagnóstico, epidemiología y control. *Phytoma-España*, No. 102, (October 1998), pp. 177-179, ISSN 1131-8988
- Penyalver, R., García, A., Ferrer, A., Bertolini, E. And López, M.M., 2000. Detection of *Pseudomonas savastanoi* pv. *savastanoi* in olive plants by enrichment and PCR. *Appl. Environ. Microbiol.*, 66, 2673–2677.
- Penyalver, R., García, A., Ferrer, A., Bertolini, E., Quesada, J.M., Salcedo, C.I., Piquer, J., Pérez-Panadés, J., Carbonell, E.A., Del Río, C., Caballero, J.M. and López, M.M., (2006) Factors affecting *Pseudomonas savastanoi* pv. *savastanoi* plant inoculations and their use for evaluation of Olive cultivar susceptibility. *Phytopathology*, 96: 313–319.
- Penyalver, R., García, A., Pérez-Panadés, J., Del Río, C., Caballero, J.M., Pinochet, J., Piquer, J., Carbonell, E., & López, M.M. (2005). Resistencia y susceptibilidad a la tuberculosis, In: Variedades de Olivo en España, Rallo, L., Barranco, D., Caballero, J.M., Del Río, C., Martín, A., Tous, J., & Trujillo, I., pp. 339-346, Junta de Andalucía, MAPA and Ediciones Mundial Prensa, ISBN 848474146X, Madrid, Spain.
- Pereira P.S., 2015. *Xylella fastidiosa* – a new menace for Portuguese agriculture and forestry. *Revista de Ciências Agrárias*, 38(2): 149-154.
- Perez J.A., J. M. Hernández, J. M. Trujillo, and H. López, "Iridoids and secoiridoids from Oleaceae," *Studies in Natural Products Chemistry*, vol. 32, pp. 303–363, 2005.
- Perez-Martinez I, Rodriguez-Moreno L, Matas IM & Ramos C (2007) Strain selection and improvement of gene transfer for genetic manipulation of *Pseudomonas*

- savastanoi* isolated from olive knots. Res Microbiol 158: 60–69.

Perez-Martinez I., Zhao Y., Murillo J., Sundin GW & Ramos C (2008) Global genomic analysis of *Pseudomonas savastanoi* pv. *savastanoi* plasmids. J Bacteriol 190: 625–635.

Pérez-Martínez I., Zhao Y., Murillo J., Sundin G.W., Ramos C., 2008. Global genomic analysis of *Pseudomonas savastanoi* pv. *savastanoi* plasmids. Journal of Bacteriology 190: 625–635.

Pérez-Mendoza, D., Aragón, I.M., Prada-Ramírez, H.A., Romero-Jiménez, L., Ramos, C., Gallegos, M.-T. and Sanjuan, J. (2014) Responses to elevated c-di-GMP levels in mutualistic and pathogenic plant-interacting bacteria. PLoS One, 9, e91645.

Pesante, A. 1938. On an unknown disease of olive. Bollettino della Stazione di Patalogia Vegetale, Roma 18:401-428.

Petri L, 1915. *Le Malattie dell'Olivo*. Florence, Italy: Istituto Micrografico Italiano.

Petri, L. (1909). Ricerche Sopra i Batteri Intestinali della Mosca Olearia. Roma: Memorie della Regia Stazione di Patologia Vegetale di Roma (in Italian).

Pierez-Martinez I., Rodriguez-Moreno L., Lambertsen L., Matas IM, Murillo J., Tegli S., Jimenez AJ & Ramos C (2010) Fate of a *Pseudomonas savastanoi* pv. *savastanoi* type III secretion system mutant in olive plants (*Olea europaea* L.). Appl Environ Microbiol 76: 3611–3619.

Potere O., Susca L., Loconsole G., Saponari M., Boschia D., Savino V. and Martelli G.P., 2015. Survey for the presence of *Xylella fastidiosa* subsp. *pauca* (strain CoDiRO) in some forestry and ornamental species in the Salento peninsula. Journal of Plant Pathology, 97 (2) : 373-376.

Proetta, U. (1995). Le malattie dell' olivo. Informatore Fitopatologico, No. 12, pp. 16-26, ISSN 0020-0735.

Purcell A.H., 2013. Paradigms: Examples from the bacterium *Xylella fastidiosa*. Annual Review of Phytopathology 51: 229- 356.

Purcell, A.H. and D.L. Hopkins. 1996. Fastidious xylem-limited bacterial plant pathogens. Annu.

Pyrowolakis E, Weltzein HC (1974) Studies on the distribution of olive knot, induced by *Pseudomonas savastanoi* (SM.) Stev. in the Greek island of Crete. Phytopathologia Mediterranea 13, 118–120.

Pyrowolakis E., Weltzien H.C., 1974. Studies on the distribution of Olive knot, induced by *Pseudomonas savastanoi* (Sm.) Stev. in the Greek island of Crete / Studi sulla distribuzione della rogna dell' Olivo da *Pseudomonas savastanoi* (Sm.) Stev. nell'isola greca di Creta. Phytopathologia Mediterranea. 13 (1/2): 118-120.

Pyrowolakis, E., and H. C. Welzien. 1974. Studies on the distribution of olive-knot, induced by *Pseudomonas*

*savastunoi* (Sm.) Stev. in the Greek island of Crete. Phytopathol. Mediterr. 13:11&120.

Quesada JM, Penyalver R, Pérez-Panadés J, Salcedo CI, Carbonell EA, López MM. 2010a. Comparison of chemical treatments for reducing epiphytic *Pseudomonas savastanoi* populations and for improving subsequent control of olive knot disease. Crop Protection 29: 1413–1420.

Quesada, J. M., Penyalver, R., and López, M. M. (2012). “Epidemiology and control of plant diseases caused by phytopathogenic bacteria: the case of olive knot disease caused by *Pseudomonas savastanoi*” in Plant Pathology, ed. C. J. Cumagun (INTECH Open Access Publisher), 299–326.

Quesada, J. M., Penyalver, R., Pérez-Panades, J., Salcedo, C. I., Carbonell, E. A., and Lopez, M. M. (2010b). Dissemination of *Pseudomonas savastanoi* pv. *savastanoi* populations and subsequent appearance of olive knot disease. Plant Pathol. 59, 262–269.

Quesada, J.M., García, A., Bertolini, E., López, M.M., & Penyalver, R. (2007). Recovery of *Pseudomonas savastanoi* pv. *savastanoi* from symptomless shoots of naturally infected olive trees. International Microbiology, Vol. 10, No. 2, (June 2007), pp. 77-84, ISSN 1139-6709.

Queseda J.M., Garcia A., Bertolini E., Lopez M.M., Penyalver R., 2007. Recovery of *Pseudomonas savastanoi* pv. *savastanoi* from symptomless shoots of naturally infected olive trees. International Microbiology 10: 77-84.

Ramos C, Matas IM, Bardaji L, Aragon IM & Murillo J (2012). *Pseudomonas savastanoi* pv. *savastanoi*: some like it knot. Mol Plant Pathol 13: 998–1009.

Redak, R. A., Purcell, A. H., Lopes, J. R. S., Blua, M. J., Mizell, R. F., and Andersen, P. C. 2004. The biology of xylem fluid-feeding insect vectors of *Xylella fastidiosa* and their relation to disease epidemiology. Annu. Rev. Entomol. 49:243-270

Rei, F.T., Henriques, M.I.C., Leitao, F.A., Serrano, J.F. and Potes, M.F. 1993. Immunodiagnosis of Cucumber mosaic cucumovirus in different olive cultivars. Bulletin OEPP/EPPO Bulletin, 23: 501-504.

Rhouma, A. Triki, M.A. Msallek, M. First report of olive anthracnose caused by *Colletotrichum gloeosporioides* in Tunisia. Phytopathol. Mediterr. 49: 95-98 (2010)

Rice R. E., Phillips P. A., Stewart-Leslie J., Sibbert G. S. (2003). Olive fruit fly populations measured in Central and Southern California. Calif. Agric. 57:122-127.

Rieuf P. Teasca G. 1970. Parasites et saprophytes des plantes au Maroc. Les cahiers de la recherche agronomique Direction de la recherche agronomique, Rabat. Marot 9.309.

Roberto, F. F., and T. Kosuge. 1987. Phytohormone metabolism in *Pseudomonas syringae* subsp. *savastanoi*,

- p. 371-380. In J. E. Fox and M. Jacobs (ed.), Molecular biology of plant growth control. Alan R. Liss, Inc., New York.
- Roberto, F. F., Klee, H., White, F., Nordeen, R., Kosuge, T. 1990. Expression and fine structure of the gene encoding NE-indole-3-acetyl-L-Iysine synthetase from *Pseudomonas savastanoi*. Proc. Natl. Acad. Sci. USA 87:734–38.
- Roca L.F., Moral J., Viruega J.R., Ávila A., Oliveira R., Trapero A., 2007. Copper fungicides in the control of olive diseases. FAO olive network. 48-50.
- Rodríguez-Moreno L., Barceló-Muñoz A., Ramos C., 2008. *In vitro* analysis of the interaction of *Pseudomonas savastanoi* pvs *savastanoi* and *nerii* with micropropagated olive plants. Phytopathology 98: 8.
- Rodríguez-Moreno, L., Barceló-Munoz, A. and Ramos, C. (2008) *In vitro* analysis of the interaction of *Pseudomonas savastanoi* pvs. *savastanoi* and *nerii* with micropropagated olive plants. Phytopathology, 98, 815–822.
- Rodríguez-Moreno, L., Jimenez, A. J., and Ramos, C. (2009). Endopathogenic lifestyle of *Pseudomonas savastanoi* pv. *savastanoi* in olive knots. Microb. Biotechnol. 2, 476–488.
- Rodriguez-Palenzuela P, Matas IM, Murillo J. (2010) Annotation and overview of the *Pseudomonas savastanoi* pv. *savastanoi* NCPPB 3335 draft genome reveals the virulence gene complement of a tumour-inducing pathogen of woody hosts. Environ Microbiol 12: 1604–1620.
- Rodríguez-Palenzuela, P., Matas, I. M., Murillo, J., Lopez-Solanilla, E., Bardaji, L., Pérez-Martínez, I., et al. (2010). Annotation and overview of the *Pseudomonas savastanoi* pv. *savastanoi* NCPPB 3335 draft genome reveals the virulence gene complement of a tumour-inducing pathogen of woody hosts. Environ. Microbiol. 12, 1604–1620.
- Rojas, A.M. (1999). Análisis fenotípicos, genéticos y filogenéticos de flora endofítica asociada a *Pseudomonas savastanoi*. Thesis (Ph. D.). Universidad San Pablo CEU, Madrid, Spain.
- Rojas, A.M., García de los Ríos, J.E., Fischer-Le Saux, M., Jiménez, P., Reche, P., Bonneau, S., Sutra, L., Mathieu-Daudé, F., & McClland, M. (2004). *Erwinia toletana* sp. nov., associated with *Pseudomonas savastanoi*-induced tree knots. International Journal of Systematic and Evolutionary Microbiology, Vol. 54, No. 6, (November 2004), pp. 2217- 2222, ISSN 1466 5026.
- Rokni-Zadeh, H., Khavazi, K., Asgharzadeh, A., Hosseini-Mazinani, M., & De Mot, R. (2008). Biocontrol of *Pseudomonas savastanoi*, causative agent of olive knot disease: antagonistic potential of non-pathogenic rhizosphere isolates of fluorescent *Pseudomonas*. Communications in Agricultural and Applied Biological Sciences, Vol. 73, No. 1, pp. 199-203, ISSN 1379 1176.
- Roussos P.A., Pontikis C.A., *In vitro* propagation of olive (*Olea europaea* L.) cv. *Koroneiki*, Plant Growth Reg. (2002) (in press).
- Roussos, P. A.; Pontikis, C. A.; Tsantili, E. Root promoting compounds detected in olive knot extract as response to infection by the bacterium *Pseudomonas savastanoi* pv *saVastanoi*. *Plant Sci.* **2002**, *163*, 533-541.
- Ryan D. and K. Robards, "Phenolic compounds in olives," *Analyst*, vol. 123, no. 5, pp. 31R–44R, 1998.
- Saad A.T., Hanna L., 2002. Two new hosts of *Pseudomonas savastanoi* and variability in strains isolated from different hosts. *Phytopathology* **92**: S71.
- Saad A.T., Melkonian S., 1992. The distribution and host range of *Pseudomonas syringae* ssp. *savastanoi* in Lebanon. *MPU/EPPO Conference on Olive Diseases, Sounion 1992*: 67.
- Saad A.T., Nienhaus F., 1969. Plant Diseases in Lebanon. Zeitschrift für Pflanzenkrankheiten (Pflanzenpathologie) und Pflanzenschutz, 76 (9/10): 537-551.
- Sabanadzovic S., Abou-Ghanem N., La Notte P., Savino V., Scarito G., Martelli G.P., 1999. Partial molecular characterization and RT-PCR detection of a putative closterovirus associated with olive leaf yellowing. *Journal of Plant Pathology* **81**: 37-45.
- Sampson P.J. and Walker J., 1982. An annotated list of plant diseases in Tasmania. Dept. of Agriculture, p. 118-121.
- Sands D.C., Schroth M.N., Hildebrand D.C., 1970. Taxonomy of Phytopathogenic Pseudomonadst. *Journal of Bacteriology*, 101 (1): 9-23.
- Sanei, S. J. and Razavi, S. E. Survey of Olive Fungal Disease in North of Iran. Annual Review & Research in Biology, 2(1): 27-36 (2012)
- Sanei, S.J. Okhovvat, S.M. Hedjaroude, Gh.A. Saremi, H. Javan-Nikkhah, M. Peltate trichome distribution and scab resistant in some olive cultivars. 3rd Biological Compound and Optimize Use of Pesticides in Agriculture, 2004, 592.
- Saponari M, Castellano MA, Grieco F, Savino V, Martelli GP. (2004) Further studies on Olive leaf yellowing-associated virus. *J Plant Pathol* 86:295–340.
- Saponari M., Alkowni R., Grieco F., Pantaleo V., Savino V., Martelli G. P., Driouech N., Hassan M., Di Terlizzi B., Digiaro M., 2002a. Detection of olive-infecting viruses in the Mediterranean Basin. *Acta Horticulturae* **586**: 787-790.
- Tsuji M., Ohta K., Tanaka K., Takikawa Y., 2017. Comparison among Japanese isolates of *Pseudomonas savastanoi* pv. *savastanoi*, causal agent of olive knot disease. *J Gen Plant Pathol*. 83:152–161.

- Saponari M., Boscia D., Altamura G., D'Attoma G., Cavalieri V., Zicca S., Morelli M., Tavano D., 2016. Pilot project on *Xylella fastidiosa* to reduce risk assessment uncertainties. Institute for Sustainable Plant Protection, National Research Council of Italy, CNR. 60pp.
- Saponari M., Mondelli G., Savino V., Martelli G.P., 2002b. Nuove acquisizioni sui virus dell'olivo. In: *Proceedings Convegno Internazionale Olivicoltura, Spoleto 2002*, 472-476.
- Saponari M., Savino V. and Martelli G.P. (2002). Trasmissione per seme di virus dell'olivo. Frut-ticoltura. 64(4), 103-105.
- Saponari M., Savino V., 2003. Virus and virus-like agents in olive, Informatore Fitopatologico 12: 26-29.
- Saponari M., Savino V., 2003. Viruses and phytoplasmas of olive [*Olea europaea* L.]. <http://agris.fao.org/agris-search/search.do?recordID=IT2004062070>
- Saponari M., Loconsole G., Cornara D., Yokomi R.K., De Stradis A., Boscia D., Bosco D., Martelli G.P., Krugner R., and Porcelli F., 2014. Infectivity and Transmission of *Xylella fastidiosa* by *Philaenus spumarius* (Hemiptera: Aphrophoridae) in Apulia, Italy. 107 (4) : 1316-1319.
- Saponari M.; Boscia D., Nigro F., Martelli G. P., 2013. Identification of DNA sequences related to *Xylella fastidiosa* in oleander, almond and olive trees exhibiting leaf scorch symptoms in Apulia (southern Italy). Journal of Plant Pathology. 95(3) : 668.
- Saponari, M., Boscia, D., Loconsole, G., Palmisano, F., Savino, V. N., Potere, O., & Martelli, G. P. (2014c). New hosts of *Xylella fastidiosa* strain CoDiRO in Apulia. Journal of Plant Pathology, 96, 611.
- Saponari, M., El Beaino, T., Grieco, F., Savino, V., & Martelli, G. P. (2001). Diagnosi di virus dell'olivo mediante l'impiego degli RNA a doppia elica. Atti Progetto POM A, 32, 793–797.
- Saponari, M., Loconsole, G., Almeida, R., Coletta-Filho, H., Martelli, G. P., & Boscia, D. (2014a). Isolation, genotyping and preliminary data on the pathogenicity of *Xylella fastidiosa* CoDiRO strain. International Symposium on the European Outbreak of *Xylella fastidiosa* in Olive, Journal of Plant Pathology, 96, S4.103.
- Sarwar M., "The theatrical usefulness of olive *Olea europaea* L.(Oleaceae family) nutrition in human health: a review," Sky Journal of Medicinal Plant Research, vol. 2, no. 1, pp. 1–4, 2013.
- Savastano L (1878) Il bacillo della tubercolosi dell'olivo Rendicontazione dell'Accademia dei Lincei 5: 2 sem., fasc. 3.
- Savastano L, 1887. Tubercolosi, iperplasie e tumori dell'olivo. Annali della Regia Scuola Superiore di Agraria di Portici 5, 1-117.
- Savastano L, 1889. Il bacillo della tubercolosi dell'olivo. Rendiconti della Regia Accademia dei Lincei 5, 92-4.
- Savastano, L. (1886). Les maladies de l'olivier, et la tuberculose en particulier. C. R. Séance. Acad. Agric. Fr. 103, 1144.
- Savastano, L., 1908, Sulla transmissibilità del bacillo della tubercolosi dell'Olivo nell'Oleandro, Bol. Arb. Ital., 4: 86-87. 2.
- Savino V. and D. Gallitelli, 1981. Cherry leaf roll in olive. Phytopathologia Mediterranea 22, 76-77.
- Savino V. and D. Gallitelli, 1983. Isolation of Cucumber mosaic virus from Olive in Italy. Phytopathologia Mediterranea 22, 76-77.
- Savino V., Baldoni L., Faggioli F., Loconsole G., Saponari M., 2011. Innovazioni per la certificazione delle produzioni vivaistiche di olivo: il contributo del progetto OLIVIVA. Acta Italus Hortus 1: 345-348.
- Savino V., D. Gallitelli and M. Barba, 1983. Olive latent ringspot virus, a newly recognized virus infecting olive in Italy. Annals of Applied Biology 103, 243–249.
- Savino V., Gallitelli D., Barba M., (1983) Olive latent ringspot virus, a newly recognized virus infecting olive in Italy. *Annals of Applied Biology* **103**, 243-249.
- Savino V., M. Barba, D. Gallitelli and G.P. Martelli, 1979. Two nepoviruses isolated from olive in Italy. Phytopathologia Mediterranea 18, 135–142.
- Savino V., Piazzola P., Di Franco A., and G.P., Martelli, 1984. Olive latent virus-2, a newly recognized virus with differently shake particles. In: Proceeding of the 6<sup>th</sup> Congress of the Mediterranean Phytopathological Union, Cairo, Egypt, 1984, 24–26.
- Savino V., Saponari M., Bottalico G., D'Ascenzo D., Barbone S., Laurentiis De G., 2004. Olive clonal and phytosanitary selection in Abruzzo region. <http://agris.fao.org/agris-search/search.do?recordID=IT2006602326>.
- Schroth MN (1995) 'Integrated control of olive knot disease.' Final report for UC Davis IPM Competitive Research Grants Program. Available online at <http://www.ipm.ucdavis.edu/calludt.cgi/> Grants Summary, Project=93FE027&PLIST=02BU007, 9 3FE027, 88BC001 (verified 18 May 2004).
- Schroth MN , Hildebrand DC 1968, A chemotherapeutic treatment for selectively eradicating crown gall and olive knot neoplasms. 58(6):848-854.
- Schroth MN, Hildebrand DC, O'Reilly HJ (1968) Off-flavor of olives from trees with olive knot tumors. *Phytopathology* **58**, 524–525.
- Schroth MN, Osgood JW, Miller TD (1973) Quantitative assessment of the effect of the olive knot disease on olive yield and quality. *Phytopathology* **63**, 1064–1065.
- Scorticini M, Rossi MP and Salerno M (2004) Relationship of genetic structure of *Pseudomonas*

- savastanoi* pv. *savastanoi* populations from Italian olive trees and patterns of host genetic diversity. Plant Pathology 53: 491–497.
- Scroth MN, 1995. *Integrated control of the olive knot disease. Annual report for 1995 – UC IPM Competitive Grants Program.* California, USA: University of California Davis. [http://ipm.ucdavis.edu/IPMPROJECT/1995/applied.html#end].
- Serçe, Ç. U.; Yalçın, S.; Gazel, M.; Çağlayan, K.; Faggioli, F., 2007. First report of *Olive latent virus 1* from olive trees in Turkey. Journal of Plant Pathology 89 (3): 73 pp.
- Sergeeva, V. Tesoriero, L. Spooner-Hart, R. Nair, N.G0. First report of *Macrophomina phaseolina* on olives (*Olea europaea*) in Australia. Plant Pathology, 2005, 34, 273-274.
- Servi D., Baştas K.K., 2012. Determination and Identification of Olive Knot Disease (*Pseudomonas savastanoi* pv. *savastanoi*) in Aydın Province. Selçuk Tarım ve Gıda Bilimleri Dergisi 26 (3): 1-8.
- Shoemaker C.A., Huffaker C.B., AND Kennett C.E., 1979. A Systems Approach to the Integrated Management of a Complex of Olive Pests. Environ. Entomol. 8: 182—189.
- Sibbett G.S., L. Ferguson, and M. Lindstrand, Olive Production Manual, University of California, Department of Agriculture and Natural Resources, 2005.
- Simpson T (2015) Utopians v. Pro!teers? Diffusion of *Xylella* in Italian olive trees. <http://xylellacodiro.blogspot.fr/2015/04/utopians-v-pro-teers.html>
- Sisto A, Cantore P, Iacobellis N (2001) Preliminary results on the response of olive cultivars to artificial inoculation with *Pseudomonas syringae* subsp. *savastanoi*. In 'Proceedings of the 11th Congress of the Mediterranean Phytopathological Union and 3rd Congress of the Sociedade Portuguesa de Fitopatologia'. Evora, Portugal, 17–20 September 2001. pp. 240–242.
- Sisto A, Iacobellis NS (1999) Olive knot disease: pathogenic and epidemiological aspects and defence strategies. *Olivo and Olio.* 2, 32–38.
- Sisto A., Cipriani M.G., Morea M., 2004. Knot formation caused by *Pseudomonas syringae* subsp. *savastanoi* on olive plants is hrp-dependent. Phytopathology 94: 484-489.
- Sisto A., Cipriani M.G., Morea M., Lonigro S.L., Valerio F., Lavermicocca P., 2010. An Rhs-like genetic element is involved in bacteriocin production by *Pseudomonas savastanoi* pv. *savastanoi*. Antonie van Leeuwenhoek, 98:505–517.
- Sisto A., Cipriani M.G., Tegli S., Cerboneschi M., Stea G., Santilli E., 2007. Genetic characterization by fluorescent AFLP of *Pseudomonas savastanoi* strains isolated from different host species. Plant pathology, 56 (3): 366–372.
- Sisto A., Morea M., Baruzzi F., Palumbo G., 2002. Differentiation of *Pseudomonas syringae* subsp. *savastanoi* strains isolated from various host plants by restriction fragment length polymorphism. Phytopathol. Mediterr. 41, 63–71
- Škarica, B., Žužić, I., Bonifačić, M. 1996. Maslina i maslinovo ulje visoke kakvoće u Hrvatskoj. Tipograf d.d., Rijeka.
- Smidt, M. L., and A. K. Vidaver. 1986. Isolation and characterization of syringacin W-1, a bacteriocin produced by *Pseudomonas syringae* pv. *syringae*. Can. J. Microbiol. 32:231–236.
- Smidt, M., and T. Kosuge. 1978. The role of indole-3-acetic acid accumulation by alpha methyl tryptophan-resistant mutants of *Pseudomonas savastanoi* in gall formation on oleanders. Physiol. Plant Pathol. 13:203–214.
- Smith ER and Rorer JB (1904). The olive tubercle. Science 480 (XIX): 416–417.
- Smith, E, D, 1908. Recent studies on the olive tubercle organism. Bull. Bur. Plant Ind. U. S. Dep. Agric. 131:2543.
- Smith, Pathogenicity of the olive knot organism on hosts related to the olive. Phytopathology 12:271--278. 1922.
- Soby S., Kirckpatrick B., Kosuge T., 1933. Characterization of an insertion sequence (IS53) located within IS51 on the iaa-containing plasmid of *Pseudomonas syringae* pv. *savastanoi*. Plasmid. 29: 135-141.
- Spahiu E., Veizi A., 2012. The Sanitation of Olive Tree Varieties from Viral Infections Using the "in vitro" Technique. Journal of International Environmental Application and Science; Konya 7.5 (Dec 2012): 901-906.
- Spooner-Hart R., Tesoriero L., Hall B., 2007. Field Guide to Olive Pests, Diseases and Disorders in Australia. Rural Industries Research and Development Corporation. 07/153.
- Surico G (1977) Histological observations on tumours of olive knot. *Phytopathologia Mediterranea* **16**, 109–125.
- Surico G and Marchi G (2003) Olive knot disease: new insights in the ecology, physiology and epidemiology of *Pseudomonas savastanoi* pv. *savastanoi*. In: Iacobellis NS, Collmer A, Utcheson SW, Mansfield JW, Morris CE, Murillo J, Schaad NW, Stead DE, Surico G and Ullrich MS (eds), *Pseudomonas syringae* and related pathogens: Biology and Genetic (pp 17–28) Kluwer Academic Publishers, Dordrecht, The Netherlands.
- Surico G., Iacobellis N.S., Sisto S., 1985a. Studies on the role of indole-3-acetic acid and cytokinins in the

- formation of knots on olive and oleander plants by *Pseudomonas syringae* pv. *savastanoi*. *Physiological Plant Pathology* 26: 309-320.
- Surico G., Sparapano L., Lerario P., Durbin R.D., Iacobellis N., 1975. Cytokinin-like activity in extracts from culture filtrates of *Pseudomonas savastanoi*. *Experientia*, 31(8):929–930.
- Surico, G. & Iacobellis, N.S. (1991) Phytohormones and olive knot disease. In Molecular Signals in PlantMicrobe Communications (ed. Verma, D.P.S.), pp. 209-227. CRC Press, Boca Raton (USA).
- Surico, G. (1993). Scanning electron microscopy of olive and oleander leaves colonized by *Pseudomonas syringae* subsp.*savastanoi*. *J. Phytopathol.* 138, 31–40.
- Surico, G., and Iacobellis, N. S. (1992). “Phytohormones and olive knot disease,” in Molecular Signals in Plant-Microbe Communications, ed. D. P. Verma (Boca Raton, FL: CRC Press), 209–227.
- Surico, G., and Lavermicocca, P. (1989). A semiselective medium for the isolation of *Pseudomonas syringae* pv. *savastanoi*. *Phytopathology* 79, 185–190. doi: 10.1094/phyto-79-185.
- Surico, G., Comai, L. and Kosuge, T., 1984. Pathogenicity of strains of *Pseudomonas syringae* pv. *savastanoi* and their indolacetic acid deficient mutants on olive and oleander. *Physiol. Biochem.*, 74, 490–493.
- Surico, G., Evidente, A., Taclobellis, N.S & Randazzo, G. (1985b) A cytokinin from the culture filtrate of *Pseudomonas syringae* pv. *savastanoi*. *Phytochemistry* 24, 1499-1 502.
- Sutic, D. and Dowson, W. J., 1963. The reaction of olive oleander and ash, cross inoculated with some strains and forms of *Pseudomonas savastanoi* (Smith ) stevens . *Phytopath. z.*, 46, 305 – 314.
- Tatlı B. ve Benlioğlu K., 2004 Aydın ve Muğla İllerinde Zeytin Ağacı Uru *Pseudomonas savastanoi* pv. *savastanoi* Hastalığı Üzerine Çalışmalar. T.C. Adnan Menderes Üniversitesi, Fen Bilimleri Enstitüsü, Yüksek Lisans Tezi, Bitki Koruma Anabilim Dalı, 62 s.
- Tegli S., Cerboneschi M., Marsili I.L., Santilli E., 2010. Development of a versatile tool for the simultaneous differential detection of *Pseudomonas savastanoi* pathovars by End Point and Real-Time PCR. 10 (156): 1-14.
- Tehabsim, A., Khadair And Jd. Janse 1991. Oeeurenee and distribution of *Pseudomonas syringae* subsp. *savastanoi* in Jordan. *Phytopatologia Mediterranea*, 30: 1,64-66p.
- Temsah M., Hanna L. and Saad A.T., 2007c. Anatomical observations of *Pseudomonas savastanoi* on *Rhamnus alaternus*. *For. Path.* 37: 64–72.
- Temsah, M., Hanna, L., & Saad, A.T. (2007a) Anatomical observations of *Pseudomonas savastanoi* on *Rhamnus alaternus*. *Forest Pathology*, Vol. 37, No. 1, (February 2007), pp. 64–72, ISSN 1439-0329.
- Temsah, M., Hanna, L., & Saad, A.T. (2007b) Histological pathogenesis of *Pseudomonas savastanoi* on *Myrtus communis*. *Journal of Plant Pathology*, Vol. 89, No. 2, (July 2007), pp. 241–249, ISSN: 1125-4653.
- Temsah, M., Hanna, L., and Saad, A. T. (2008). Anatomical pathogenesis of *Pseudomonas savastanoi* on olive and genesis of knots. *J. Plant Pathol.* 90, 225– 232.
- Teviotdale BL (1994) Diseases of olive, In ‘Olive production manual’. (Eds L Ferguson, GS Sibbett, GC Martin) p. 107. University of California Publication No. 3353.
- Teviotdale BL, Krueger B, 1998 olive knot control and assessment of copper resistance in *Pseudomonas savastanoi*. In: Stapleton JJ, Summers CG, Teviotdale BL, Goodell PB, Prather TS (Eds) *Plant Protection Quarterly* (Vol. 8), Kearney Plant Protection Group, pp 1-4.
- Teviotdale, B. L. 2005. Diseases of olive. Pages 119-121 in: Olive Production Manual, 2nd ed. G. S. Sibbett and L. Ferguson. eds. University of California Agricultural and Natural Resources, Publ. 3353, Oakland, CA.
- Teviotdale, B. L., and Krueger, W. H. 2004. Effects of timing of copper sprays, defoliation, rainfall, and inoculum concentration on incidence of olive knot disease. *Plant Dis.* 88:131-135.
- Tsuji M, Ohta K, Tanaka K, Takikawa Y (2015). First report of knot disease on olive (*Olea europaea*) in Japan caused by *Pseudomonas savastanoi* pv. *savastanoi*. *Plant Dis* 99:1445
- Tjamos, E. C.; Graniti, A.; Smith, I. M.; Lamberti, F.; Eds.; Conference on olive diseases. *EPPO Bull.* **1993**, 23, 365-550.
- Tommasini R., 1992. Degree thesis (Tesi di Laurea) Universita` degli Studi, Florence, Italy.
- Tous J., Romero A., Hermoso J.J., 2010. New trends in olive orchard design for continuous mechanical harvesting. *Adv. Hort. Sci.*, 24(1): 43-52
- Trapero. A, Roca. L. F., Moral, J., López-Escudero, F. J., Blanco-López, M. A. (2009). Enfermedades del olivo. *Phytoma España: La Revista Profesional de Sanidad Vegetal*, 209, 18-28 (in Spanish).
- Trigui M., Ben Hsouna A., Hammami I., Culoli G., Ksantini M., Tounsi S., Jaoua S., 2013. Efficacy of *Lawsonia inermis* leaves extract and its phenolic compounds against olive knot and crown gall diseases. *Crop Protection* 45: 83-88.
- Úrbez-Torres, J. R., Peduto, F., Vossen, P. M., Krueger, W. H., and Gubler, W. D. 2013. Olive twig and branch dieback: Etiology, incidence, and distribution in California. *Plant Dis.* 97:231-244.

- USDA, "National Genetic Resources Program," Germplasm Resources Information Network—(GRIN), 2003.
- Varanda C, Nolasco G, Clara M, Félix M (2014a) Genetic diversity of the coat protein gene of Olive latent virus 1 isolates. *Arch Virol* 159: 1351–1357.
- Varanda C.M.R., Santos S., Clara M.I.E., Félix R., 2015a. Olive mild mosaic virus transmission by *Olpidium virulentus*. *European Journal of Plant Pathology* 142 (1): 197–201.
- Varanda CMR, Machado M, Martel P, Nolasco G, Clara MIE, et al. (2014b) Genetic Diversity of the Coat Protein of Olive Mild Mosaic Virus (OMMV) and Tobacco Necrosis Virus D (TNV-D) Isolates and Its Structural Implications. *PLoS ONE* 9(10): e110941. doi:10.1371/journal.pone.0110941
- Varanda CMR, Machado M, Martel P, Nolasco G, Clara MIE, Félix MR (2014) Genetic Diversity of the Coat Protein of Olive Mild Mosaic Virus (OMMV) and Tobacco Necrosis Virus D (TNV-D) Isolates and Its Structural Implications. *PLoS ONE* 9(10): e110941. doi:10.1371/journal.pone.0110941. pmid:25350108.
- Varanda, C. M. R., Cardoso, J. M. S., Félix, M. R. F., Oliveira, S., & Clara, M. I. E. (2010). Multiplex RT-PCR for detection and identification of three necroviruses that infect olive trees. *European Journal of Plant Pathology*, 127, 161–164.
- Varanda, C.M.R., Félix, M.R.F., & Clara, M.I.E. (2008). DsRNA analysis and RT-PCR assays to evaluate viral infections: the case of necroviruses in olive. *Journal of Plant Pathology*, 90 (2): 385.
- Varanda, C. M. R., Félix, M. R., Soares, C. M., Oliveira, S., & Clara, M. I. E. (2011a). Specific amino acids of Olive mild mosaic virus coat protein are involved on transmission by *Olpidium brassicae*. *Journal of General Virology*, 92, 2209–2213.
- Varanda, C. M. R., Silva, M. S. M. R., Félix, M. R., & Clara, M. I. E. (2011b). Evidence of Olive mild mosaic virus transmission by *Olpidium brassicae*. *European Journal of Plant Pathology*, 130, 165–172.
- Varanda, C.M.R., Cardoso, J.M.S., Félix, M.R.F., Oliveira, S., & Clara, M.I.E (2009). Evaluation of necroviruses occurrence by specific RT-PCR in olive. 4th European Meeting of the IOBC/WPRS working Group Integrated Protection of Olive Crops, Cordoba, Spain, Book of Abstracts, p 104.
- Varanda, C.M.R., 2011. Molecular characterization of *Olea europaea* L. necrovirus transmission by fungal vectors. Doctoral thesis. <http://hdl.handle.net/10174/9134>
- Varvaro L, Surico G (1978a) Comportamento di diverse cultivars d'olivo (*Olea europaea* L.) alla inoculazione artificiale con *Pseudomonas savastanoi* (E.F. Smith) Stevens. *Phytopathol Mediterr* 17:174–177.
- Varvaro L, Surico G (1978b) Moltiplicazione di *Pseudomonas savastanoi* (E.F. Smith) Stevens nei tessuti dell'olivo (*Olea europaea* L.). *Phytopathol Mediterr* 17:179–186
- Varvaro L. and Martella L., 1993. Virulent and avirulent isolates of *Pseudomonas syringae* subsp. *savastanoi* as colonizers of olive leaves: evaluation of possible biological control of the olive knot pathogen. *Bulletin* 23, 423-427.
- Varvaro, L. & Ferrulli, M. (1983) [Survival of *Pseudomonas syringae* pv. *saw.stanoi* on the leaves of two olive cultivars.] *Phytopathologia Mediterranea* 22, 1-4 (in Italian).
- Varvaro, L. & Surico, G. (1984) Epiphytic survival of wild types of *Pseudomonas syringae* pv. *saoastanoi* and their Iaa- mutants on olive leaves. In Proceedings of the 2nd Working Group on *Pseudomonas syringae* pathouars, pp. 20-22. Sounion (GR).
- Poggi Pollini C., Bissani R., Giunchedi L., Vindimian E., 1996. First Report of Phytoplasma Infection in Olive Trees (*Olea Europea* L.). *Journal of Phytopathology*. 144 (2): 109–111.
- Pasquini G., Marzachì C., Poggi Pollini C., Faggioli F., Ragozzino A., Bissani R., Vischi A., Barba M., Giunchedi L. and Boccardo G., 2000. *Journal of Plant Pathology*, 82 (3): 213-219.
- Danielli A., Bertaccini A., Vibio M. and Repetti S. (1996). Identificazione di fitoplasmi associati a scopazzi dell'olivo. *Atti Convegno Annuale della Società Italiana di Patologia Vegetale*, Udine, Italy .28-29.
- Martelli G.P., 1999. Infectious diseases and certification of olive: an overview. *EPPO Bulletin*. 29 (1-2) : 127–133.
- Spooner-Hart R., Tesoriero L., Hall B., 2007. Field Guide to Olive Pests, Diseases and Disorders in Australia. Rural Industries Research and Development Corporation. 68p
- Ahangaran A., Khezri S., Habibi M.K., Alizadeh A., Mohammadi G.M., 2006. The first report of detection of a phytoplasma in olive trees in a botanic collection in Iran. *Communications in Agricultural and Applied Biological Sciences*. 71(3):1133-1138.
- Font I., Abad P., Dally E.I., Davis R.E. and Jordà C. (1998). Nueva enfermedad en el olivar español. *Phytoma España*. 102, 211-212.
- Marzachi C., Poggi Pollini C., 2000. Suggestion of a detection protocol for phytoplasma identification in olive trees [*Olea europaea* L. - Italy]. Istituto di Servizi per il Mercato Agricolo Alimentare. <http://agris.fao.org/agris-search/search.do?recordID=IT2001062319>.
- Serrone P. del, Faggioli F., Barba M., Arzone A., Tarquini A., 1996. Phytoplasmas associated to olive witches'-broom. <http://agris.fao.org/agris-search/search.do?recordID=IT9760533>.

- Marzachì C., Veratti F., d'Aquilio M., Vischi A., Conti M. and Boccardo G., Molecular hybridization and PCR amplification of nonribosomal DNA to detect and differentiate stolbur phytoplasma isolates from Italy. *Journal of Plant Pathology* 82 (3): 201-212.
- Ghayeb Zamharir M., Razavi R., 2016. First finding of a 'Candidatus Phytoplasma fraxini'-related strain associated with disease of olive in Iran. *New Disease Reports* (2016) 34, 10.
- Varvaro, L. & Surico, G. (1987) Multiplication of wild types of *Pseudomonas syringae* pv. *savastanoi* and their indoleaceatic acid-deficient mutants in olive tissues. In *Plant Pathogenic Bacteria* (eds Civerolo, E.L. et al.), pp. 556~565. Martinus Nijhoff, Dordrecht (NL).
- Vovlas C., Parrella G., Di Franco A., 2002. Infekzioni naturali del virus latente dell'olivo su piante di ricino in Grecia. *Informatore Fitopatologico*, 7-8:66-68.
- Wallander E., V.A., Albert, "Phylogeny and classification of Oleaceae based on rps16 and trnL-F sequence data," *American Journal of Botany*, vol. 87, no. 12, pp. 1827–1841, 2000.
- Warcup JH, Talbot PHB (1981) 'Host-pathogen index of plant diseases in South Australia.' (Waite Agricultural Research Institute, University of Adelaide).
- Wells, J.M., Casano, F.J. and Surico, G., 1991. Fatty acid composition of *Pseudomonas syringae* pv. *savastanoi*. *J. Phytopathol.*, 133, 152–162.
- Weltzien H., 1963. Plant Diseases in the Middle East. Observations in the Republic of Lebanon, *Phytopathologia Mediterranea*. 2(4): 260-264.
- White, S.M.; Bullock, J.M.; Hooftman, D.A.P; Chapman, D.S.. 2014 Modelling the spread of *Xylella fastidiosa* in Puglia, Italy. [Lecture] In: *International Symposium on the European Outbreak of Xylella fastidiosa in Olive, Gallipoli, Italy, 21-24 Oct 2014*. (Unpublished)
- Wilson E.E., 1935. The olive knot disease: Its conception, development, and control. *Hilgardia IX* 4: 233-257.
- Wilson J.A., Stephenson B.P., Gill G.S.C., Randall J.L. and Vieglais C.M.C., 2004. Principles of response to detections of new plant pest species and the effectiveness of surveillance. *Biosecurity*. 156-160.
- Wilson, E. E. and Magie, A., 1963. Physiological, serological, and pathological evidence that *Pseudomonas tonelliana* is identical to *Pseudomonas savastanoi*. *Phytopathology* . 53, 6, 653 – 659.
- Wilson, E. E., and Maggie, A. R. 1964. Systemic invasion of the host plant by the tumor-inducing bacterium, *Pseudomonas savastanoi*. *Phytopathology* 54:576-579.
- Wilson, E. E., Heskett, M. G., Johnson, M. L., and Kosuge, T., 1972. Metabolic behavior of *Pseudomonas savastanoi* isolates from olive and oleander on certain carbohydrate and amino substrates. *Phytobathiology* , 62, 3, 349 – 355.
- Wilson, E.E. (1965). Pathological histogenesis in oleander tumors induced by *Pseudomonas savastanoi*. *Phytopathology*, Vol. 55, pp. 1244-1249, ISSN 0031-949X.
- Wilson, E.E., & Ogawa, J.M. (1979). Fungal, bacterial, and certain nonparasitic diseases of fruit and nut crops in California, Division of Agricultural Sciences, University of California, ISBN 093187629X, Berkeley, USA.
- Wilson, M., & Lindow, S.E. (1992). Relationship of total viable and culturable cells in epiphytic populations of *Pseudomonas syringae*. *Applied and Environmental Microbiology*, Vol. 58, No. 12, (December 1992), pp. 3908-3913, ISSN 0099-2240.
- Wong F., Cooksey D.A., Costa H.S., 2004. Documentation and characterization of *Xylella fastidiosa* strains in landscape hosts. CDFA Pierce's Disease Control Program Progress Report. Proceedings Pierce's Disease Research Symposium San Diego, USA: 121-123.
- Yalçın S., 2008. RT-PCR detection and characterisation of virus spread in olive orchards in the east mediterranean region. University, Graduate School of Natural and Applied Sciences, Hatay (Turkey)Mustafa Kemal Univ., Faculty of Agriculture, Hatay (Turkey). Div. of Plant Protection.
- Yamada T., Lee P.D., Kosuge T., 1986. Insertion sequence elements of *Pseudomonas savastanoi* nucleotid sequence and homology with *Agrobacterium tumefaciens* transfer DNA. *Proc. Natl. Acad. Sci. USA* 83: 8263-8267.
- Yaseen T., Drago S., Valentini F., Elbeaino T., Stampone G., Digiaro M. and D'onghia A.M., 2015. On-site detection of *Xylella fastidiosa* in host plants and in "spy insects" using the real-time loop-mediated isothermal amplification method. *Phytopathologia Mediterranea*, 54(3) : 488–496.
- Young J.M., Wilkie J.P., Fletcher M.J., Park D., Pennycook S.R., Triggs C.M., Richard D., Watson W., 2004b. Relative tolerance of nine olive cultivars to *Pseudomonas savastanoi* causing bacterial knot disease. *Phytopathol. Mediterr.* 43, 395–402.
- Young JM (1987) Orchard management and bacterial diseases of stonefruit. *New Zealand Journal of Experimental Agriculture* 15, 257–266.
- Young JM, Saddler GS, Takikawa Y, Deboer SH, Vauterin L, Gardan L, Gvozdyak RI, Stead DE, 1996. Names of plant pathogenic bacteria 1864–1995. ISPP Subcommittee on Taxonomy of Phytopathogenic Bacteria. *Review of Plant Pathology* 75, 721–63.
- Young JM, Triggs CM (1994) Evaluation of determinative tests for pathovars of *Pseudomonas syringae* van Hall 1902. *The Journal of Applied Bacteriology* 77, 195–207.

- Young, J. M. (2004). Olive knot and its pathogens. *Australas. Plant Pathol.* 33, 33–39.
- Youssef, S. A., Moawed, S. M., El-Sayed, M., & Shalaby, A. A. (2010). Detection of olive tree viruses in Egypt by onestep RT-PCR. 21st International Conference on Virus and other Graft Transmissible Diseases of Fruit Crops, Julius-Kühn-Archiv, 427, 51–55.
- Zohary D., M. Hopf, and E. Weiss, Domestication of Plants in the Old World: The Origin and Spread of Domesticated Plants in Southwest Asia, Europe, and the Mediterranean Basin, Oxford University Press, Oxford, UK, 2012.
- Žužić, I., Ciglar, I. 1987. Usmjerena i integrirana zaštita masline. A. G. Matoš, Samobor.