

Evaluation of Storage Capacity of Iron Fortified Yogurt by Physico-chemical, Chemical and Microbiological Analysis

Debasmita De¹, Dr. BinataNayak²

¹ Junior Research Fellow, Department of Home Science (Food & Nutrition Division) University of Calcutta, Kolkata, West Bengal, India

² Assistant Professor, Department of Home Science (Food & Nutrition Division) University of Calcutta, Kolkata-700027, West Bengal, India

Abstract—Yogurt has gained widespread consumer acceptance. It is excellent source of calcium and protein and other nutrients but it contains very little iron. In this study yogurt was fortified with ammonium ferrous sulfate in three different concentrations (20mg, 30mg, 40mg/kg iron). Yogurt samples were analyzed physico-chemically, chemically and microbiologically at 1st, 3rd, 5th day of storage. Physicochemical and chemical result shows that there was significant difference between storage period and different sample concentration. Iron in the fortified samples had no significant effect in *Lactobacillus* count. The results suggest possibility of making good quality yogurt by fortifying milk with ammonium ferrous sulfate.

Keywords—Ammonium ferrous sulphate, Chemical, fortification, Microbiological, Physicochemical, Yogurt.

I. INTRODUCTION

Anaemia is a most common world-wide problem in the young children, pregnant woman and adolescent girl. Nutritional anaemia may be defined as the condition that results from the inability of the erythropoietic tissues to maintain a normal haemoglobin concentration on account of inadequate supply of one or more essential nutrients leading to reduction in the total circulating haemoglobin.[1] Most of the anaemias are due to inadequate supply of nutrients like iron, folic acid and vitamin B12, proteins, amino acids, vitamins A, C, and other vitamins of B-complex group i.e., niacin and pantothenic acid are also involved in the maintenance of haemoglobin level.[2]

Globally, anaemia affects 1.62 billion people, which corresponds to 24.8% of the population. The highest prevalence is in preschool-age children (47.4%), and the lowest prevalence is in men (12.7%). However, the population group with the greatest number of individuals affected is pregnant women (41.8%) [3]. In women, anaemia may become the underlying cause of maternal mortality and perinatal mortality. Nearly 50 per cent of

women of reproductive age and 26 per cent of men in the age group of 15-59 years are anaemic[4].

To reduce the prevalence rate of anaemia three types of measures are taken: 1. Dietary improvement, 2. Supplementation, 3. Food fortification.[1],[5],[6] The iron found in food can be highly bioavailable as in the case with heme iron which is found in red meat. The iron present in other products of vegetable origin contain non-heme iron has disadvantage of interacting with substance in food that inhibits absorption such as tannin and phytates[7]. The best way to prevent problems associated with iron deficiency is through iron fortification of food for whole population or certain group [8]. Yogurt has gained widespread consumer acceptance. It is excellent source of calcium and protein, but it contains very little iron. Therefore dairy products are good for iron fortification because they have high nutritive value reach target population and are widely consumed[9],[10],[11]. The ideal iron compound used as fortificants should supply high bioavailability iron, it should not affect the nutritional value or sensory properties of food [12],[13],[14],[15].

The purpose of this study is to prepare iron-fortified yogurt with ammonium ferrous sulfate at three different concentration (20mg, 30mg, 40mg/kg milk) as it covers respectively 9.52%, 14.28% and 19.04% of RDA of iron of an adult woman. And yogurt samples were analyzed chemically and microbiologically during 1st, 3rd, 5th day of storage.

II. MATERIALS & METHODS

The study was designed to prepare Iron fortified yogurt and physicochemical, chemical, microbial analysis of iron fortified yogurt at 1st, 3rd, 5th day of storage period. Methodologies adopted for this analysis:

1. Preparation of iron fortified yogurt.[6]
2. Physicochemical analysis of iron fortified yogurt
 - i) Whey separation by centrifugation method. [16]
 - ii) Volume of supernatant by syneresis index. [16]

iii) Determination of Total Titratable Acidity .[17]

iv) Moisture content determination. [18]

v) Total solid content determination. [18]

3. Chemical analysis of iron fortified yogurt.

i) Determination of protein by Lowry method. [19]

ii) Determination of iron by Wong's method. [20]

.4. Microbial analysis of iron fortified yogurt.

i) Enumeration of Lactobacillus count. [21]

Preparation of Iron fortified Yogurt: Locally available AmulTaja toned homogenized pasteurized milk was taken. Milk was fortified with ammonium ferrous sulfate. The milk was divided into four portions. The first portion was not fortified with iron and regarded as control. The rest three portions were fortified with ammonium ferrous sulfate in different concentration respectively 20mg, 30mg, 40mg iron/kg milk. Then milk was inoculated with yogurt culture and filled into plastic cups, covered and kept at room temperature until a firm curd was formed (approximately 6-7 hours). The resultant yogurt was kept in a refrigerator for 5 days at 4°C.[6]

Statistical analysis: This was done by Two way Analysis of Variance (first factor storage period, second factor sample concentration).

III. RESULT & DISCUSSION

3.1 Physicochemical analysis

3.1.1 Volume of whey

Sample	1st	3rd	5th	F ratio	P value
Yogurt (NF)	0.592 ±0.001	0.672 ±0.009	0.866 ±0.02	Between columns=34.85	0.00084
Yogurt (20mg)	0.694 ±0.009	0.86 ±0.06	0.91 ±0.03	Between rows=404.10	0.0005
Yogurt (30mg)	1.293 ±0.05	1.399 ±0.02	1.496 ±0.02		
Yogurt (40mg)	1.475 ±0.01	1.576 ±0.04	1.636 ±0.01		

Inference: Two Way ANOVA shows that there was a significant difference between columns and rows. So null hypothesis is rejected and alternative hypothesis is accepted.

3.1.2 Volume of supernatant by syneresis index (%)

Sample	1st	3rd	5th	F ratio	P value
Yogurt (NF)	11.85 ±0.03	13.44 ±0.19	17.32 ±0.48	Between columns=38.74	0.00075
Yogurt (20mg)	13.88 ±0.19	17.2 ±1.2	18.2 ±0.66	Between rows=430.30	0.0005

Yogurt (30mg)	25.86 ±1.03	27.98 ±0.48	29.92 ±0.53		
Yogurt (40mg)	29.2 ±0.5	31.52 ±0.81	32.72 ±0.3		

Inference: Two Way ANOVA shows that there was a significant difference between columns and rows. So null hypothesis is rejected and alternative hypothesis is accepted

3.1.3 Total Titratable Acidity (%):

Sample	1st	3rd	5th	F ratio	P value
Yogurt (NF)	0.27 ±0.005	0.29 ±0.005	0.32 ±0.01	Between columns=19.90	0.0021
Yogurt (20mg)	0.27 ±0	0.28 ±0.005	0.29 ±0.005	Between rows=6.18	0.03538
Yogurt (30mg)	0.27 ±0.005	0.30 ±0.005	0.31 ±0.005		
Yogurt (40mg)	0.28 ±0.005	0.32 ±0.01	0.34 ±0.01		

Inference: Two Way ANOVA shows that there was a significant difference between columns and rows. So null hypothesis is rejected and alternative hypothesis is accepted

3.1.4 Moisture(%):

Sample	1st	3rd	5th	F ratio	P value
Yogurt (NF)	86 ±0.7	87.2 ±0.5	88.8 ±1	Between columns=27.06	0.00119
Yogurt (20mg)	87.4 ±0.5	88.8 ±0.27	89.14 ±0.91	Between rows=9.08	0.0158
Yogurt (30mg)	87.6 ±0.9	88.4 ±0.7	89.2 ±0.63		
Yogurt (40mg)	88 ±0.7	89 ±1.7	89.4 ±0.77		

Inference: Two Way ANOVA shows that there was a significant difference between columns and rows. So null hypothesis is rejected and alternative hypothesis is accepted

3.1.5 Total solid(%):

Sample	1st	3rd	5th	F ratio	P value
Yogurt (NF)	14 ±0.7	12.8 ±0.5	11.2 ±1	Between columns=30.09	0.00101
Yogurt (20mg)	12.6 ±0.5	11.2 ±0.27	10.86 ±0.91	Between rows=9.04	0.01593
Yogurt (30mg)	12.4 ±0.9	11.6 ±0.7	10.8 ±0.63		

Yogurt (40mg)	12 ±0.7	11 ±1.7	10.6 ±0.77		
---------------	------------	------------	---------------	--	--

Inference: Two Way ANOVA shows that there was a significant difference between columns and rows. So null hypothesis is rejected and alternative hypothesis is accepted

3.2 Chemical analysis:

3.2.1 Protein (mg/ml):

Sample	1st	3rd	5th	F ratio	P value
Yogurt (NF)	31 ±0.5	31.3 ±0.9	31.4 ±1.2	Between columns=26.99	0.00119
Yogurt (20mg)	31.5 ±0.5	32 ±1	32.3 ±0.7	Between rows=80.98	0.00054
Yogurt (30mg)	32 ±2	32.4 ±0.6	32.5 ±0.5		
Yogurt (40mg)	32.2 ±0.11	32.44 ±0.55	32.56 ±1.05		

Inference: Two Way ANOVA shows that there was a significant difference between columns and rows. So null hypothesis is rejected and alternative hypothesis is accepted

3.2.2 Iron (µg/ml)

Sample	1st	3rd	5th	F ratio	P value
Yogurt (NF)	4 ±0.5	3.5 ±0.2	2.5 ±0.4	Between columns=35.80	0.00081
Yogurt (20mg)	21.06 ±0.40	20.03 ±1.15	19.5 ±0.8	Between rows=13149.72	0.0005
Yogurt (30mg)	31.5 ±0.8	31 ±0.5	30.5 ±0.5		
Yogurt (40mg)	43.5 ±0.5	42.2 ±0.75	41.53 ±0.95		

Inference: Two Way ANOVA shows that there was a significant difference between columns and rows. So null hypothesis is rejected and alternative hypothesis is accepted

3.3 Microbial analysis

3.3.1 Lactobacillus count (cfu/ml) 10^{10}

Sample	1st	3rd	5th	F ratio	P value
Yogurt (NF)	10.2 ±0.3	9.6 ±0.2	9.9 ±0.1	Between columns=1.13	0.40785
Yogurt (20mg)	11.8 ±0.8	11.03 ±0.55	11.3 ±0.3	Between rows=4.44	0.06589
Yogurt (30mg)	9 ±0.5	9.5 ±0.5	11.03 ±1.05		
Yogurt (40mg)	11 ±0.1	10.3 ±0.5	10.7 ±0.3		

Inference: Two Way ANOVA shows that there was a non-significant difference between columns and rows. So null hypothesis is accepted.

IV. CONCLUSION

Yogurt is a most important health beneficial nutritious probiotic. It is a product of the lactic acid fermentation of milk. Result shows that during storage period volume of syneresis increased in all samples but this increase is significant in non-fortified sample. There was a steady increase in Total titratable acidity during storage period. The pH was decreasing due to accumulation of lactic acid as a bacterial culture was breaking down lactose in order to obtain energy. This observation is in agreement with the previous study by Nkhata et al[22]. The metabolic enzymatic activity of the yogurt starter culture could be the reason for increases in acidity which could be responsible for decreasing lactobacillus spp. count although statistical analysis shows there was nonsignificant difference in lactobacillus count during storage period and sample concentration. Statistical analysis of moisture, total solid, iron and protein shows that there was a significant difference between sample concentration and storage period. Present study shows all fortified yogurt samples were nutritionally rich and acceptable, suggesting that yogurt is a suitable vehicle for iron fortification.

REFERENCES

- [1] Srilakshmi. B, "Nutrition Science", 3rd Ed. Chapter 10. pp-164-180. ISBN 10: 81-224-2147-4 New Age International (P) Limited. 2008.
- [2] Lee GR; Herbert V. In: Lukens J, Paraskevas P, Greer JP, Rodgers GM, editors. "Wintrobe's clinical hematology". Baltimore, Maryland USA: William & Wilkins, pp 228-266. 1998,
- [3] WHO, WHO global database on anaemia / Edited by Bruno de Benoist, Erin McLean, Ines Egli and Mary Cogswell, 2008. http://whqlibdoc.who.int/publications/2008/9789241596657_eng.pdf retrieved on July February 6th, 2014
- [4] Ezzati M; Lopus AD; Dogers A; Vander HS; Murray C, *Lancet*, "A Cross-Sectional Study of Hemoglobin Disorders in Pregnant Women Attending Two Urban Hospitals in Eastern Coast of Odisha, India". 360, pp1347-1360, 2002.
- [5] Nayak. Band. Madhavan KNair. Department of Biophysics NINICMR Hyderabad. "In Vitro Bioavailability of Iron from Wheat Flour Fortified with Ascorbic acid, EDTA and SHMP with or without Iron. "Food Chemistry Volume 80, Issue 4, pp.545-550. April 2003

- [6] Amira M. El-Kholy, Osman .M, Gouda.A and Wafaa A. Ghareeb , "Fortification of Yoghurt with Iron". World Journal of Dairy & Food Sciences 6 (2): pp159-165, 2011 ISSN 1817-308X IDOSI Publications, 2011
- [7] Van den Broek N, Letsky EA. "Etiology of anaemia in pregnancy in south Malawi". Am L Clin Nutr. 72 (1Suppl): 247S-256S, 2000.
- [8] Biswas.T and Nayak .B, "Iron fortification of Green Gram Dhal flour and its iron invitro bioavailability as a preventive measure of anaemia", American International Journal of Research in Formal, Applied and natural science, 9(1), pp. 64-66, 2014.
- [9] Sadler A.M., Lacroix D.E., Alford J.A, "Iron content of Baker's and cottage cheese made from fortified skim milks". J. Dairy Sci., 56: 1267-1270, 1973.
- [10] Tamime A. Y., Robinson R. K, "Yoghurt: Science and Technology". 2nd ed. CRC Press, Boca Raton, FL. 1999
- [11] Tamime A. Y., Deeth H. C. "Yogurt: technology and biochemistry", Journal of Food Protection 43 939–977, 1980.
- [12] United States Department of Agriculture. USDA Specifications for Yogurt, Non -fat Yogurt and Low fat Yogurt, Dairy Programs. 2001.
- [13] Lee W. J., Lucey J. A, "Formation and Physical Properties of Yogurt", Asian-Aust. J. Anim. Sci. Vol. 23, No. 9 : 1127 – 1136, 2010.
- [14] Zhang D., Mahoney A.W, "Iron fortification of process Cheddar cheese". J. Dairy Sci., 74: 353-358, 1991.
- [15] Saha.S, Dr. Nayak .B, "Evaluation of Organoleptic Properties and Storage Capacity of Green Gram Dhal Flour (*Vigna Radiata*) Fortified with Iron and Its Absorption Promoters ", AIJRFANS 14(1), March-May, 2016, pp. 26-29
- [16] Guzman- Gonzalez M., Morais F. and Amigo L, "Influence of skimmed milk concentrate replacement by dairy products in a low-fat set type yoghurt model system. Use of caseinates, coprecipitate and blended dairy powder", J. Sci. Food Agri, 80: 433-438.
- [17] Behrad S., Yusof M.Y., Goh K. L. and Baba A. S, "Manipulation of Probiotics Fermentation of Yogurt by Cinnamon and Licorice: Effects on Yogurt Formation and Inhibition of Helicobacter Pylori Growth in vitro". World Academy of Science, Engineering and Technology, 2009; 36: 12-29.
- [18] AOAC, 1995, "Official Methods of Analysis of the Association of Official Analytical Chemists", Vol III 16th ed., AOAC Arlington, VA.
- [19] Lowry, O.H., N.J. Rosenbrough, A.L. Farr and R.L. Randall, "Protein measurement with the Folin Phenol Reagent". J. Biol. Chem., 193: 265-275. 1951.
- [20] N. Raghuramulu, KM Nair and S. Kalyanasundaram. NIN. ICMR. "A Manual of Laboratory Technique" 77. Page 409. 1982
- [21] Dr. Pande Sunil S and Gupta Priyanka, "A Promising Functional Dairy Food Using Probiotics With Kalmegh ". Rajiv Gandhi Biotechnology Centre, Rashtrasant Tukdoji Maharaj Nagpur University, L.I.T. Premises, Nagpur-440 033 (M.S.), India. ISSN 2349-8870 Volume: 2 Issue: 3 376-384 Year: 2015.
- [22] Nkhata Gilliard Smith , Ustunol Zeynep, Menevseoglu Ahmed, "Iron Fortification Yogurt and Pasteurised Milk". Journal of nutritional Health and Food Science. Pub: April 13, 201