

First record of the Pacific bluefin tuna *Thunnus orientalis* (Temminck & Schlegel, 1844) from the coast off Sur, Sultanate of Oman

Shama Zaki, Juma Al-Mamary, Abdul Aziz Al-Marzouqi, Lubna Al-Kharusi

Marine Science and Fisheries Centre, Ministry of Agriculture and Fisheries, Muscat, Sultanate of Oman

Abstract— A single specimen of the Pacific bluefin tuna *Thunnus orientalis* was caught on 11 May 2017 in a long-line operated about 40 nautical miles off the coast of Sur atconfluent of Sea of Oman and Arabian Sea coast of Oman. This first record of its occurrence indicates the extension of distributional range of the species to the Arabian Sea coast of Oman.

Keywords— coast off Sur, Pacific Bluefin, Arabian Sea.

I. INTRODUCTION

Oman has a very long coastline of 3165 km with connections to three seas namely the Arabian Gulf, the Gulf of Oman and the Arabian Sea. The country has rich fish biodiversity (Al-Jufaily *et al.*, 2010). The biodiversity of fish fauna of Omani coasts had been reported by several earlier workers (Boulenger, 1887; Steindachner, 1902; Regan, 1905; Norman, 1939; White & Barwani, 1971; Randall, 1995; Al-Abdessalaam, 1995; Fouda *et al.*, 1997). In addition, there is scope for inclusion of several new species and new records to the list of known species of Omani fishes (McKoy *et al.*, 2009; Jawad & Al-Mamry, 2009). As the result, a number of new records of fish species were reported from the Sea of Oman and Arabian Sea off Oman from recent studies (Jawad & Al-Mamry, 2009; Jayabalan *et al.*, 2010; Al-Jufaily *et al.*, 2010; Jawad *et al.* 2014).

Fishes of the genus *Thunnus*(Perciformes: Scombridae) are represented by eight species namely albacore, *T.*

alalunga (Bonnaterre, 1788), southern bluefin tuna, *T. maccoyii* (Castelnau, 1872), bigeye tuna, *T. obesus* (Lowe, 1839), Pacific bluefin tuna, *T. orientalis* (Temminck & Schlegel, 1844), Atlantic bluefin tuna, *T. thynnus* (Linnaeus, 1758), yellowfin tuna, *T. albacares* (Bonnaterre, 1788), blackfin tuna, *T. atlanticus* (Lesson, 1831) and longtail tuna, *T. tonggol* (Bleeker, 1851) from world oceans (Froese & Pauly, 2009. Godsil, & Byers, 1945; Collette & Nauen 1983). Of these, Pacific bluefin tuna, *T. orientalis* is a commercially valuable species and is widely distributed in the Pacific Ocean (Froese & Pauly, 2009) and seasonally inhabiting subarctic, temperate, and tropical waters in the North Pacific Ocean as well as temperate waters in the Southern Hemisphere around Australia and New Zealand (Collette *et al.*, 2014).

Pacific bluefin tuna grows to a maximum weight of ~650 kg, a total length of ~300 cm and age of at least 20 years and it is the second largest species of tuna, after the Atlantic bluefin tuna (*Thunnus thynnus*) (Collette and Nauen, 1983; Kitagawa *et al.*, 2007; Collette *et al.*, 2014). Pacific blue fin tuna is highly prized in the Japanese sashimi markets and during 2014 about 17,076 t of fish were landed; however, *T. orientalis* is reported as 'Vulnerable' under IUCN red list of species (Collette *et al.*, 2014).

(Collette and Nauen, 1983; Foreman and Ishizuka, 1990; Bayliff, 1994)



Fig.1: Map of Sultanate of Oman

From the Omani waters so far only three species such as *T. tonggol*, *T. albacares* and *T. obesus* are known (Godsil & Byers, 1945; Collette *et al.*, 2001; Randall, 1995; Froese & Pauly, 2011). In the present report, the occurrence of single specimen of Pacific Bluefin tuna *T. orientalis* off the coast of Suratconfluent of Sea of Oman and Arabian Sea coast of Oman (Fig.1) is recorded for the first time.

SYSTEMATICS

Order: PERCIFORMES

Family: SCOMBRIDAE

Genus: *Thunnus*

Species: *orientalis*



Fig.2: Specimen of *T. orientalis* collected from Oman

II. MATERIAL EXAMINED

One specimen, Fig. 2, is 250 cm total length (TL), (204 cm standard length- SL; 224 cm fork length- FL); caught by long line; 11 May 2017. Atconfluent of Sea of Omanand Arabian Sea coast of Oman. It is difficult to say precisely because the fisherman did not have GPS (global positioning system).

The sample was brought to the Marine science laboratory at the Marine Science and Fisheries Centre, Ministry of Agriculture and Fisheries Wealth, Sultanate of Oman for analysis. After weighing,the morphometric and meristic characters of the fish were recorded following standard procedures (John& Schaefer, 1949). Then, the fish was cut

open to identify the sex, maturity stage of the gonad and feeding intensity. The liver and air bladder structure were observed. Otoliths were extracted from head for subsequent age analysis. The muscle tissue was collected for genetic analysis. The fish sample was buried to preserve and display the bones in the museum.

Description

The fish weighed 237 kg and was a mature male. The feeding intensity was 1/2full stomach and the food was in fully digested state. Ventral surface of liver was striated with blood vessels. Air bladder was irregular. There were 36 gill rakers in the first left gill arch (Fig. 3)



Fig.3: First gill arch- Total gill rakers- 36.

The morphometric characters (Table- 1) and meristic counts (Table- 2) of the specimen agree with the descriptions of the species *T. orientalis* which was further confirmed by the genetic analysis of mitochondrial DNA with D-Loop (NCBI. Accession No. JN631213.1).

Body: Fusiform, almost round, very robust in front and tapering towards caudal peduncle. Head lengthless than body depth. Mouth large,teeth small and conical in a single series. The first dorsal height less (10.7% of FL) than the second dorsal fin height (16.1% of FL) in the present specimen; pectoral fin short and less than head length and about 17% of FL. In *T. orientalis*, the short pectoral fin and heights of 1st and 2nd dorsal fins are considered as the

prominent morphometric characters to identify the species (Nelson, 2006).

Coloration: Lower side of belly was silvery white but no other colour pattern was visible. This might be due to the delay in bringing the specimen to the laboratory after three days. However, in fresh specimens lower sides and belly are with faint colourless transverse lines alternated with rows of faint colourless dots (Collette *et al.*, 1983; Nelson, 2006; Tamura & Takagi, 2009).The first and second dorsal fins were dark bluish with dusky yellow tips. Dorsal and anal Finlets were yellowish, base reddish-brown towards middle and tips. Median caudal keel was black.

Table.1: Important morphometric measurements of *T. orientalis*

S.No.	Morphometric measurements	cm	In FL (%)
1	Head length	57	25.4
2	Head depth	51	22.8
3	Pre-opercle length	44	19.6

4	Post orbital distance	33	14.7
5	Eye Diameter	6	2.7
6	Upper jaw length	20	8.9
7	Body depth	131	58.5
8	Girth	59	26.3
9	Distance snout to eye	19	8.5
10	Distance snout to nostril	17	7.6
11	Distance snout to 1 st dorsalfin base	66.5	29.7
12	Distance snout to 2 nd dorsalfin base	116	51.8
13	Distance between 1 st and 2 nd dorsalfin	22	9.8
14	1 st Dorsalfin length	49	21.9
15	1 st Dorsalfin height	24	10.7
16	2 nd Dorsalfin length	20	8.9
17	2 nd Dorsalfin height	36	16.1
18	Anal fin length	15	6.7
19	Anal fin height	36	16.1
20	Pectoralfin length	38	17.0
21	Caudal peduncle length	8	3.6
22	Caudalfin spread	71	31.7
23	Median keel height	5	2.2
24	Median keel length	25	11.2

Table.2: Meristic Counts of *T. orientalis*

S. No.	Meristic Counts	No.
1	1 st Dorsal fin spines	XIII
2	2 nd Dorsal fin rays	15
3	Dorsal Finlets	9
4	Pectoral fin rays	30
5	Anal fin rays?	12
6	Anal Finlets	8
7	Gill rakers	36
8	Keels	3

III. DISCUSSION

Thunnus orientalis is an epipelagic oceanic species that forms school by size and performs wide horizontal and vertical migrations and seasonally moves close to the shore (Magnuson, 1973; 1978; Kitagawa *et al.* 2007). This species is distributed in depth range of 1 to 550 m (Froese & Pauly, 2009; IUCN, 2014). The fish has great tolerance of sea-surface temperature ranging from 17°C to 23°C and be able to dive deep waters as cold as ~3°C (fish Base). It is a voracious predator feeding primarily on a variety of small schooling fishes such as anchovies, sardines, herrings, menhaden and mackerels and also on squids, crabs and other less sessile organisms (Collette and Nauen, 1983; Allain, 2005; Swada *et al.*, 2005).

The record of *T. orientalis* from the Arabian Sea coast of Oman is significant as the species has been found distributed in north and south Pacific (Ashida *et al.* 2015; Collette *et al.* 2014, Lewis, 2012; Itoh *et al.* 2003; Tanaka *et al.* 2006). Although, this species moves in schools, only single specimen was caught presently. This would lead to speculations such as incidental occurrence owing to disorientation or be linked to the ecological changes that occurred due to rise in water temperature in its distributional ranges causing a decline in prey organisms and hence, migrated to more abundant prey related regions or to unknown biotic and/or anthropogenic factors (Sharp & Dizon, 1978; Kimura *et al.*, 2010).

The species *T. orientalis* is represented by a single Pacific-wide stock that is found primarily in the North Pacific

Ocean (Kitigawa *et al.*, 2007). Hence, it may be presumed that the migration of the species from the north Pacific to the Arabian Sea coast of Oman might have occurred through the Bay of Bengal. Hence, it would be interesting to know whether the extension of the species distribution is very recent to the countries bordering the Bay of Bengal coast or already established populations are available in the region. Intensive research in this line is needed to better explain the occurrence of a solitary Pacific blue fin tuna from Oman. Tagging studies would reveal the possible routes of movement pattern of *T. orientalis* if any, from north Pacific to the western Indian Ocean.

ACKNOWLEDGEMENT

I would like to thank the fisherman for taking the effort to report to Ministry of Agriculture and Fisheries. Thanks are due to the Director General of Fisheries Research and Director of Marine Science and Fisheries Centre for arranging to purchase the fish sample. Thanks to all staff of the centre for their effort in handling such huge fish specimen.

REFERENCES

- [1] Al-Abdessalaam T.Z.S., 1995. Marine species of the Sultanate of Oman. Ministry of Agriculture and Fisheries, Sultanate of Oman. Publication No. 46/95, 412 pp.
- [2] Al-Jufaili, S.M., Hermosa G., Al-Shuaili S.S. and Al-Mujaini A., 2010. Oman fish biodiversity. *Journal of King Abdulaziz University: Marine Science* 21 (1): 3-51.
- [3] Allain, V., 2005. Diet of four tuna species of The Western and Central Pacific Ocean. *SPC Fisheries Newsletter* 114: 30-33.
- [4] Ashida H, Suzuki N, Tanabe T, Suzuki N, Aonuma E., 2015. Reproductive condition, batch fecundity, and spawning fraction of large Pacific bluefin tuna *Thunnus orientalis* landed at Ishigaki Island, Okinawa, Japan. *Environ Biol Fish.*, 98:1173–83.
- [5] Asiapacifish-
<http://www.asiapacifish.org/index.php/species/item/27-pacific-bluefin-tuna#biology>.
- [6] Boulenger G.A. (1887) An account of the fishes obtained by Surgeon-Major A.S.G. Jayakar at Muscat, east coast of Arabia. *Proceedings of the Zoological Society of London* 1889, 653–667.
- [7] Collette, B.B. and Nauen.C.E. 1983. FAO Species Catalogue Vol.2 Scombrids of the World: An annotated and illustrated catalogue of tunas, mackerels, bonitos, and related species known to date. Retrieved from Rome, Italy.
- [8] Collette, B., Reeb, C and Block, B.A., 2001. Systematics of the tunas and mackerels (Scombridae). In: Block, B. A., & Stevens, E. D. (Eds.). pp. 1-33, Tuna: physiology, ecology, and evolution (Vol. 19). Gulf Professional Publishing.
- [9] Collette, B., Fox, W., Juan Jordan, M., Nelson, R., Pollard, D., Suzuki, N. and Teo, S. 2014. *Thunnus orientalis*. The IUCN Red List of Threatened Species 2014:170341A65166749. <http://dx.doi.org/10.2305/IUCN.UK.-3.RLTS.T170341A65166749.en>. Downloaded on 09 July 2017.
- [10] Dickson, K.A., 1995. Unique adaptations of the metabolic biochemistry of tunas and billfishes for life in the pelagic environment. *Env. Biol. Fish.*, 42: 65-97.
- [11] Fischer, W. and Bianchi, G. (ed.). 1984. FAO species identification sheets for fishery purposes, Western Indian Ocean (Fishing Area 51), Prepared and printed with the support of the Danish International Development Agency (DANIDA), FAO, Rome. Vol. 1: 6 p.
- [12] Fish catalogue-
<http://researcharchive.calacademy.org/research/ichthyology/catalog/fishcatget.asp?genid=1905>.
- [13] Fossun *et al.*, 2000. *FAO Species Identification Field Guide for Fishery Purposes*. Rome. FAO 293pp.
- [14] Fouda, M.M., Hermosa Jr. G.V. and Al-Harthi M.S., 1997. Status of fish biodiversity in the Sultanate of Oman. Convention on Biological Diversity, First National Report, Directorate General of Nature Reserves, Ministry of Regional Municipalities and Environment, Sultanate of Oman, pp. 64–104.
- [15] Froese, R. and Pauly, D., 2009. FishBase. World Wide Web) electronic publication. <http://www.fishbase.org>.
- [16] Froese R. and Pauly D. (eds), 2011. *FishBase*. World Wide Web electronic publication, www.fishbase.org. version (06/2011).
- [17] Godsil, H.C. and Byers.R.D. 1945. A Systematic Study of the Pacific Tunas. The Quarterly Review of Biology, 20(4), 390-390.
- [18] Itoh T, Tsuji, S., Nitta, A., 2003. Migration patterns of young Pacific bluefin tuna (*Thunnus orientalis*) determined with archival tags. *Fish Bull.* 101:514–34.
- [19] Jayabalan N, Shama Zaki and Lubna Al-Kharusi, 2010. First record of the slender ponyfish *Equulites elongatus* from the Arabian Sea coast of Oman. *Marine Biodiversity Records*, Vol. 3, 1-3.

- [20] Jawad L.A. and Al-Mamry, J., 2009. First record of *Antennarius coccineus* from the Gulf of Oman and second record of *Antennarius indicus* from the Arabian Sea coast of Oman. *Marine Biodiversity Records* 2, 1-3.
- [21] Jawad, L.A., Juma, M. Al-Mamry, Haithem, K., Al-Busaidi, 2014. New record of the keeltail pomfret, *Taractes rubescens* (Jordan & Evermann, 1887) (Perciformes: Bramidae) from the Sea of Oman. *International Journal of Marine Science*, Vol.4, No.25 227-230 (doi: 10.5376/ijms.2014.04.0025).
- [22] Kimura, S., Kato, Y., Kitagawa, T. and Yamaoka, N., 2010. Impacts of environmental variability and global warming scenario on Pacific bluefin tuna (*Thunnus orientalis*) spawning grounds and recruitment habitat. *Progress in Oceanography*, 86(1–2), 39-44.
- [23] Kitagawa, T., Boustany, A.M., Farwell, C.J., Williams, T.D., Castleton, M.R. and Block, B.A. 2007. Horizontal and vertical movements of juvenile bluefin tuna (*Thunnus orientalis*) in relation to seasons and oceanographic conditions in the eastern Pacific Ocean. *Fisheries Oceanography*, 16(5), 409-421.
- [24] Lewis, A.D., 2012 Conservation Management and Measures (CMM) Report on CMM 2010-04 (Pacific Bluefin Tuna). Western and Central Pacific Fisheries Commission. Northern Committee Eighth Regular Session. 3-6 September 2012. Nagasaki, Japan.
- [25] Magnuson, J.J., 1973. Comparative study of adaptations for continuous swimming and hydrostatic equilibrium of scombroid and xiphoïd fishes. *Fish Bull.*, 71 (2):337-356.
- [26] Magnuson, J.J., 1978. Locomotion by scombrid fishes: hydrodynamics, morphology, and behavior. *Fish Physiol.*, 7: 239-313.
- [27] John, C.M. and Schaefer, M.B. 1949. Definitions of body dimensions used in describing tunas. *U.S. Fish Wild. Serv. Fish. Bull.* 51:241-244.
- [28] McKoy J, Bagley N, Gauthier S and Devine J (2009) *Fish resources assessment survey of the Arabian Sea coast of Oman*, Technical Report 1, Bruce Shallard and Associates, New Zealand, 189 pp.
- [29] NCBI- <http://www.NCBI>.
- [30] Nelson, J.S., 2006. *Fishes of the world*. 4th ed. John Wiley and Sons, Hoboken, New Jersey.
- [31] Norman J.R. (1939) Fishes in the John Murray Expedition 1933–34. *British Museum (Natural History)*, *London Scientific Reports* 7, 1–116.
- [32] Randall, J. E., 1995. Coastal Fisheries of Oman, United States of America, University of Hawaii Press. 432 pp.
- [33] Regan C.T., 1905. On fishes from the Persian Gulf, the Sea of Oman, and Karachi, collected by Mr. F.W. Townsend. *Journal of the Bombay Natural History Society* 16, 318–333.
- [34] Sawada, Y, Okada, T., Miyashita, S, Murata, O. and Kumai, H., 2005. Completion of the Pacific bluefin tuna *Thunnus orientalis* (Temminck & Schlegel) life cycle. *Aquaculture Research*, 36(5), 413-421.
- [35] Sharp, G.D. and Dizon, A.D., 1978. The Physiological Ecology of Tunas. Academic Press, New-York: 485 p.
- [36] Stromme, T., 1986. Pelagic and Demersal Fish Resources of Oman, Results of the RV Fridjof Nansen Surveys in Oman 1983-1984, Institute of Marine research, Bergen, Norway.
- [37] Steindachner, F., 1902. Fische aus Su'darabien und Soko'tra. *Denkschriften der Akademie der Wissenschaften in Wien* 71, 123–168.
- [38] Tamura, Y. and Takagi, T. 2009. Morphological features and functions of bluefin tuna change with growth. *Fisheries Science*, 75(3), 567-575.
- [39] Tanaka, Y., Satoh, K., Iwahashi, M., Yamada, H., 2006. Growth-dependent recruitment of Pacific bluefin tuna *Thunnus orientalis* in the northwestern Pacific Ocean. *Mar. Ecol. Prog. Ser.*, 319:225–235.
- [40] White A.W. and Barwani M.A. (1971) Common sea fishes of the Arabian Gulf and Gulf of Oman. Dubai: Trucial States Council, 170 pp.